

# Automorphisms of Non-Singular Nilpotent Lie Algebras

Aroldo Kaplan and Alejandro Tiraboschi

Communicated by D. Müller

**Abstract.** For a real, non-singular, 2-step nilpotent Lie algebra  $\mathfrak{n}$ , the group  $\text{Aut}(\mathfrak{n})/\text{Aut}_0(\mathfrak{n})$ , where  $\text{Aut}_0(\mathfrak{n})$  is the group of automorphisms which act trivially on the center, is the direct product of a compact group with the 1-dimensional group of dilations. Maximality of some automorphisms groups of  $\mathfrak{n}$  follows and is related to how close is  $\mathfrak{n}$  to being of Heisenberg type. For example, at least when the dimension of the center is two,  $\dim \text{Aut}(\mathfrak{n})$  is maximal if and only if  $\mathfrak{n}$  is of Heisenberg type. The connection with fat distributions is discussed.

*Mathematics Subject Classification 2010:* 17B30, 16W25.

*Key Words and Phrases:* Lie groups, Lie algebras, Heisenberg type groups.

## 1. Introduction

A 2-step nilpotent real Lie algebra  $\mathfrak{n}$  with center  $\mathfrak{z}$  is called *non-singular* [E], or said *to satisfy hypothesis (H)* [M], or be the Lie algebra of a *Métivier group* [MS], if  $\text{ad } x : \mathfrak{n} \rightarrow \mathfrak{z}$  is onto for any  $x \notin \mathfrak{z}$ . Equivalently, the bracket defines a vector-valued antisymmetric form

$$[\cdot, \cdot] : \mathfrak{v} \times \mathfrak{v} \rightarrow \mathfrak{z},$$

$\mathfrak{v} = \mathfrak{n}/\mathfrak{z}$ , such that the 2-forms  $\lambda([u, v])$  on  $\mathfrak{v}$  are non-degenerate for all  $\lambda \in \mathfrak{z}^*$ ,  $\lambda \neq 0$ . Here we shall call such algebras *fat* for short, since they are the symbols of fat distributions (as opposite to "flat", or integrable, ones [Mo]), which motivate the questions.

Let  $m = \dim(\mathfrak{z})$ . While for  $m = 1$  there is only one fat algebra up to isomorphisms, for  $m \geq 2$  there is an uncountable number of isomorphism classes and for  $m \geq 3$  they form a wild set.

In this paper we study the size of groups of automorphisms of  $\mathfrak{n}$ .  $\text{Aut}(\mathfrak{n})$  itself is the semidirect product of the group  $G(\mathfrak{n})$  of graded automorphisms of  $\mathfrak{n} = \mathfrak{v} \oplus \mathfrak{z}$  with the abelian group  $\text{Hom}(\mathfrak{v}, \mathfrak{z})$ , times the group of dilations  $(t, t^2)$ . Hence, we concentrate on  $G = G(\mathfrak{n})$ .

We prove that there is an exact sequence

$$1 \rightarrow G_0 \rightarrow G \rightarrow \text{O}(m)$$

where  $G_0$  is the subgroup of  $G$  of elements that act trivially on the center. In other words, there are positive-definite inner products on  $\mathfrak{z}$  which are invariant under all of  $\text{Aut}(\mathfrak{n})$ .

If a metric  $g$  is also given on  $\mathfrak{v}$ , as in the case of the nilpotentization of a subriemannian structure, we also consider the subgroups  $K_0, K$ , of graded automorphisms that leave  $g$  invariant, which define a compatible exact sequence

$$1 \rightarrow K_0 \rightarrow K \rightarrow O(m).$$

Next, we compute the terms in this sequence and the images  $G/G_0$  and  $K/K_0$ , proving that the exactness of

$$1 \rightarrow \text{Lie}(K_0) \rightarrow \text{Lie}(K) \rightarrow \mathfrak{so}(m) \rightarrow 1$$

is equivalent to  $\mathfrak{n}$  being of Heisenberg type, while the exactness of

$$1 \rightarrow \text{Lie}(G_0) \rightarrow \text{Lie}(G) \rightarrow \mathfrak{so}(m) \rightarrow 1$$

is strictly more general. As to  $G_0(\mathfrak{n})$ , we describe it in detail for the case  $m = 2$ , leading a proof that, at least in that case,  $\dim \text{Aut}(\mathfrak{n})$  is maximal if and only if  $\mathfrak{n}$  is of Heisenberg type.

In the last section we explain the connection with the Equivalence Problem for fat subriemannian distributions.

Algebras of Heisenberg type are defined as follows [K]. If  $\mathfrak{v}$  is a real unitary module over the Clifford algebra  $\text{Cl}(\mathfrak{z})$  associated to a quadratic form on  $\mathfrak{z}$ , the identity

$$\langle z, [u, v] \rangle_{\mathfrak{z}} = \langle z \cdot u, v \rangle_{\mathfrak{v}}$$

with  $z \in \mathfrak{z} \subset \text{Cl}(\mathfrak{z})$ ,  $u, v \in \mathfrak{v}$ , defines a fat  $[\cdot, \cdot] : \mathfrak{v} \times \mathfrak{v} \rightarrow \mathfrak{z}$ . Alternatively, they are characterized by possessing a positive-definite metric such that the operator  $z \cdot$  defined by the above equation satisfies  $z \cdot (z \cdot v) = -|z|^2 v$ .

It follows from Adam’s theorem on frames on spheres [H] that for any fat algebra there is an Heisenberg type algebra with the same  $\dim \mathfrak{z}$  and  $\dim \mathfrak{v}$ . That these were, in some sense, the most symmetric, was expected from the properties of their sublaplacians [BTV] [CGN] [GV] [K], but we found no explicit statements in this regard.

Related properties of the automorphism groups of nilpotent Lie groups are studied in [P] and [MS].

## 2. Automorphisms of fat algebras

Let  $\mathfrak{n}$  be a 2-step Lie algebra with center  $\mathfrak{z}$  and let  $\mathfrak{v} = \mathfrak{n}/\mathfrak{z}$ , so that

$$\mathfrak{n} \cong \mathfrak{v} \oplus \mathfrak{z} \tag{1}$$

and the Lie algebra structure is encoded into the map

$$[\cdot, \cdot] : \Lambda^2 \mathfrak{v} \rightarrow \mathfrak{z}.$$

Let  $n = \dim \mathfrak{v}$  and  $m = \dim \mathfrak{z}$ . Relative to a basis compatible with (1), the bracket becomes an  $\mathbb{R}^m$ -valued antisymmetric form on  $\mathbb{R}^n$  and an automorphism is a matrix of the form

$$\begin{pmatrix} a & 0 \\ c & b \end{pmatrix}, \quad a \in \text{GL}(n), \quad b \in \text{GL}(m), \quad c \in \mathbb{R}^{n \times m}$$

such that

$$b([u, v]) = [au, av].$$

$\text{Aut}(\mathfrak{n})$  always contains the normal subgroup  $\mathfrak{D}(\mathfrak{n})$  of dilations and translations

$$\begin{pmatrix} tI_n & 0 \\ c & t^2I_m \end{pmatrix}, \quad t \in \mathbb{R}^*, \quad c \in \mathbb{R}^{n \times m}.$$

Let

$$G = G(\mathfrak{n}) = \left\{ \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}, \quad a \in \text{SL}(n), \quad b \in \text{GL}(m), \quad b([u, v]) = [au, av] \right\}.$$

Then  $\text{Aut}(\mathfrak{n})$  is the semidirect product of  $G(\mathfrak{n})$  with  $\mathfrak{D}(\mathfrak{n})$ . Let

$$G_0 = G_0(\mathfrak{n}) = \left\{ \begin{pmatrix} a & 0 \\ 0 & I_m \end{pmatrix}, \quad a \in \text{SL}(n), \quad [au, av] = [u, v] \right\},$$

the subgroup of automorphisms that act trivially on the center. These are Lie groups,  $G_0$  is normal in  $G$ , and the quotient group

$$G/G_0$$

can be identified with the group of  $b \in \text{GL}(\mathfrak{z})$  such that  $b([u, v]) = [au, av]$  for some  $a \in \text{SL}(\mathfrak{v})$ . Obviously,

$$\dim \text{Aut}(\mathfrak{n}) = nm + 1 + \dim(G/G_0) + \dim(G_0). \tag{2}$$

**Theorem 2.1.** *Let  $\mathfrak{n}$  be a fat algebra with center  $\mathfrak{z}$ . Then there is a positive definite metric on  $\mathfrak{z}$  invariant under  $G(\mathfrak{n})$ .*

**Proof.** Fix arbitrary positive inner products on  $\mathfrak{v}$  and  $\mathfrak{z}$ . For  $z \in \mathfrak{z}$ ,  $u, v \in \mathfrak{v}$

$$(T_z u, v)_{\mathfrak{v}} = (z, [u, v])_{\mathfrak{z}}$$

defines a linear map  $z \mapsto T_z$  from  $\mathfrak{z}$  to  $\text{End}(\mathfrak{v})$ . Clearly,

$$\mathfrak{n} \text{ fat} \Leftrightarrow T_z \in \text{GL}(\mathfrak{v}) \quad \forall z \neq 0.$$

Hence the hypothesis insures that the Pfaffian

$$P(z) = \det(T_z)$$

is non-zero on  $\mathfrak{z} \setminus \{0\}$ . This is a homogeneous polynomial of degree  $n$ , so it satisfies

$$k\|z\|^n \leq |P(z)| \leq K\|z\|^n \tag{3}$$

where  $k, K$  are the minimum and maximum values of  $|P|$  on the unit sphere, which are positive.

Let now  $g_{a,b} := \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} \in \text{Aut}(\mathfrak{n})$ . Then

$$T_{b^\dagger z} = a^\dagger T_z a$$

because  $(T_{b^\dagger z} u, v)_\mathfrak{v} = (b^\dagger z, [u, v])_\mathfrak{z} = (z, b([u, v]))_\mathfrak{z} = (z, [au, av])_\mathfrak{z} = (T_z au, av)_\mathfrak{v} = (a^\dagger T_z au, v)_\mathfrak{v}$ . Consequently

$$P(b^\dagger z) = (\det a)^2 P(z).$$

In particular, if  $g \in G$  then  $P(b^\dagger z) = P(z)$ . This implies

$$k \|b^\dagger z\|^n \leq |P(b^\dagger z)| = |P(z)| \leq K \|z\|^n$$

for all  $z$ , therefore  $\|b\| \leq \sqrt[n]{K/k}$ . The group of  $b \in \text{GL}(\mathfrak{z})$  such that  $g_{a,b} \in \text{Aut}(\mathfrak{n})$  for some  $a \in \text{SL}(\mathfrak{v})$ , is therefore bounded in  $\text{End}(\mathbb{R}^m)$ . Its closure is a compact Lie subgroup of  $\text{GL}(\mathfrak{z})$ , necessarily contained in  $\text{O}(\mathfrak{z})$  for some positive definite metric. ■

From now on  $\mathfrak{z}$  will be assumed endowed with such invariant metric. If a metric  $g$  on  $\mathfrak{v}$  is also fixed, as in the case of the nilpotentization of a subriemannian structure, define the groups

$$K = K(\mathfrak{n}, g) = \left\{ \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}, a \in \text{SO}(\mathfrak{v}), b \in \text{O}(\mathfrak{z}), [au, av] = b[u, v] \right\}$$

$$K_0 = K_0(\mathfrak{n}, g) = \left\{ \begin{pmatrix} a & 0 \\ 0 & I \end{pmatrix}, a \in \text{SO}(\mathfrak{v}), [au, av] = [u, v] \right\}.$$

Let  $\mathfrak{g}, \mathfrak{g}_0, \mathfrak{k}, \mathfrak{k}_0$  be the Lie algebras of  $G, G_0, K, K_0$  respectively. Then there is the commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathfrak{g}_0 & \rightarrow & \mathfrak{g} & \rightarrow & \mathfrak{so}(m) \\ & & \uparrow & & \uparrow & & \uparrow \\ 0 & \rightarrow & \mathfrak{k}_0 & \rightarrow & \mathfrak{k} & \rightarrow & \mathfrak{so}(m) \end{array}$$

where the vertical arrows are the inclusions. Below we prove that the bottom sequence extends to

$$0 \rightarrow \mathfrak{k}_0 \rightarrow \mathfrak{k} \rightarrow \mathfrak{so}(m) \rightarrow 0$$

if and only if  $\mathfrak{n}$  is of Heisenberg type. This is not the case for the top one: the condition that

$$0 \rightarrow \mathfrak{g}_0 \rightarrow \mathfrak{g} \rightarrow \mathfrak{so}(m) \rightarrow 0$$

is exact defines a class of fat algebras strictly larger than Heisenberg type algebras. We describe it in the next section for  $m = 2$ .

**Proposition 2.2.** *Let  $\mathfrak{n} = \mathfrak{v} + \mathfrak{z}$  be an algebra of Heisenberg type. There is a metric on  $\mathfrak{z}$  such that  $\mathfrak{g}/\mathfrak{g}_0 \cong \mathfrak{so}(m)$ .*

**Proof.** There is an inner product in  $\mathfrak{v}$  such that the  $J_i = T_i$ 's satisfy the Canonical Anticommutation Relations

$$J_w J_z + J_z J_w = -2 \langle z, w \rangle I.$$

For  $\|z\| = 1$  let  $r_z \in O(\mathfrak{z})$  be the reflection through the hyperplane orthogonal to  $z$  and  $J_z \in SL(\mathfrak{v})$  be as above. Then

$$g_{(J_z, -r_z)} = \begin{pmatrix} J_z & 0 \\ 0 & -r_z \end{pmatrix} \in \text{Aut}(\mathfrak{n}).$$

Indeed,

$$\begin{aligned} (w, [J_z u, J_z v]) &= (J_w J_z u, J_z v) = (-J_z J_w u - 2(z, w)u, J_z v) \\ &= -(J_z J_w u, J_z v) - 2(z, w)(u, J_z v) = (J_w u, J_z J_z v) + 2(z, w)(J_z u, v) \\ &= -(J_w u, v) + 2(z, w)(J_z u, v) = (J_{-w+2(z,w)z} u, v) \\ &= (-w + 2(z, w)z, [u, v]) = (-r_z(w), [u, v]) \\ &= (w, -r_z([u, v])), \end{aligned}$$

so that

$$-r_z([u, v]) = [J_z u, J_z v].$$

The Lie group generated by the  $-r_z$  has finite index in  $O(\mathfrak{z})$ . ■

**Corollary 2.3.** *Let  $\mathfrak{n}$  be a fat algebra with center of dimension  $m$ . Then*

$$\dim(K/K_0) \leq \dim(G/G_0) \leq m(m-1)/2$$

*with equality achieved for any Heisenberg type algebra of the same dimension with center of the same dimension.*

Since  $\text{Aut}(\mathfrak{n})/\text{Aut}_0(\mathfrak{n}) = (G/G_0) \times (\text{dilations})$ , one obtains

**Corollary 2.4.** *Let  $\mathfrak{n}$  be a fat algebra with center of dimension  $m$ . Then*

$$\dim(\text{Aut}(\mathfrak{n})/\text{Aut}_0(\mathfrak{n})) \leq 1 + m(m-1)/2,$$

*with equality achieved for any Heisenberg type algebra of the same dimension and with center of the same dimension.*

A converse for Corollary 2.3 is

**Theorem 2.5.** *If  $\mathfrak{n}$  is fat with center of dimension  $m$  and*

$$\dim(K/K_0) = m(m-1)/2$$

*for some metric on  $\mathfrak{v}$ , then  $\mathfrak{n}$  is of Heisenberg type.*

**Proof.** The hypothesis implies that  $\mathfrak{k}/\mathfrak{k}_0 = \mathfrak{g}/\mathfrak{g}_0 \cong \mathfrak{so}(m)$ , so that  $K/K_0$  acts transitively among the  $|z| = 1$ . For  $\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$  in this group,  $-T_{bz} = aT_z a^{-1}$ , hence  $T_{bz}^2 = aT_z^2 a^{-1}$ . Since  $T_z$  is invertible, we can choose the metric such that  $T_{z_0}^2 = -I$  for any given  $z_0$ . Therefore  $T_z^2 = -I$  for all  $|z| = 1$ , which implies the assertion. ■

Maximal dimension means there are isomorphisms

$$\text{Lie}(K/K_0) = \text{Lie}(G/G_0) \cong \mathfrak{so}(m).$$

Therefore the simply connected covers are isomorphic:  $\text{Spin}(m) \cong \widetilde{(G/G_0)}_e$ . The induced homomorphism

$$\text{Spin}(m) \rightarrow (G/G_0)_e$$

may or may not extend to a homomorphism

$$\text{Pin}(m) \rightarrow G/G_0.$$

If it does extend, it may or may not be injective, in which case it is an isomorphism. Therefore, among the algebras for which  $\dim(G/G_0)$  is maximal, those for which  $\text{Pin}(m) \cong G/G_0$  can be regarded as the most symmetric.

**Theorem 2.6.** *Suppose  $\mathfrak{n}$  is a 2-step graded algebra such that  $\text{Aut}(\mathfrak{n})$  contains a copy of  $\text{Pin}(m)$  inducing the standard action on  $\mathfrak{z}$ . Then  $\mathfrak{n}$  is of Heisenberg type.*

**Proof.** The assumption implies that there is a linear map  $\mathfrak{z} \rightarrow \text{End}(\mathfrak{v})$ , denoted by  $z \mapsto J_z$  such that  $J_z^2 = -|z|^2 I$  for all  $z$  and

$$[J_z u, J_z v] = r_z([u, v])$$

for  $u, v \in \mathfrak{v}$ ,  $z \in \mathfrak{z}$ ,  $|z| = 1$ , where  $r_z$  is the reflection in  $\mathfrak{z}$  with respect of the line spanned by  $z$ .  $\text{Pin}(m)$  is the group generated by the  $J_z$ 's with  $\|z\| = 1$ , which acts linearly on  $\mathfrak{v}$  and is compact. Fix a metric on  $\mathfrak{v}$  invariant under it.

We get, as in the proof of Theorem 2.1, that if  $\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} \in \text{Aut}(\mathfrak{n})$ , then

$$T_{b^{\flat}z} = a^{\flat} T_z a.$$

In particular:

$$T_{r_x(z)} = J_x T_z J_x.$$

If  $x = z$ , we get  $T_z = -J_z T_z J_z$ , thus  $T_z J_z = -J_z^{-1} T_z = J_z T_z$ . If  $x \perp z$ , we get  $T_z = J_x T_z J_x$ , thus  $T_z J_x = J_x^{-1} T_z = -J_x T_z$ . It follows that  $T_z^2$  commutes with  $J_z$  and with  $J_w$ ,  $w \perp z$ .

Now, let  $z \in \mathfrak{z}$  and  $w \perp z$ . Let  $R_w(t)$  the  $2t$ -rotation from  $z$  towards  $w$ . Then  $R_w(t) = r_z r_{w(t)}$ , with  $w(t) = \cos(t)z + \sin(t)w$ . It follows that

$$\begin{pmatrix} J_z J_{w(t)} & 0 \\ 0 & R_w(t) \end{pmatrix}$$

is an orthogonal automorphism and, therefore, satisfies

$$T_{R_w(t)z} = (J_z J_{w(t)})^\dagger T_z (J_z J_{w(t)}).$$

Since  $(J_z J_{w(t)})^\dagger = (J_z J_{w(t)})^{-1}$ ,

$$T_{R_w(t)z}^2 = (J_z J_{w(t)})^\dagger T_z^2 (J_z J_{w(t)}) = J_{w(t)} J_z T_z^2 J_z J_{w(t)}.$$

Since  $T_z^2$  commutes with  $J_z$  and  $J_w$ ,

$$T_{R_w(t)z}^2 = T_z^2 J_{w(t)} J_z J_z J_{w(t)} = -T_z^2 J_{w(t)} J_{w(t)}. \tag{4}$$

But  $J_{w(t)}^2 = -I$ , so that (4) implies that

$$T_{R_w(t)z}^2 = T_z^2.$$

For all  $z' \in \mathfrak{z}$  we can choose  $w \in \mathfrak{z}, t \in \mathbb{R}$  such that  $R_w(t)z = z'$ , so we get

$$T_{z'}^2 = T_z^2, \quad \text{for all } z' \in \mathfrak{z}, |z'| = 1.$$

The antisymmetry of the bracket implies that  $T_z$  is skew-symmetric. Rescaling the scalar product on  $\mathfrak{v}$  we obtain that  $T_z^2 = -I$ , so  $T_{z'}^2 = -I$  for all  $z' \in \mathfrak{z}, |z'| = 1$ . Therefore  $\mathfrak{n}$  is of Heisenberg type. ■

### 3. The case of center of dimension 2

In this section we compute the groups  $G, G_0, G/G_0$  in the case  $m = 2$ . The various types are parametrized by pairs

$$(\mathbf{c}, \mathbf{r}) \in (\mathbb{U}^\ell / \text{SL}(2, \mathbb{R})) \times \mathbb{Z}_+^\ell$$

where  $\mathbb{U}$  is the upper-half plane and  $2\ell = 2 \sum r_j = \dim \mathfrak{n} - 2$ . As a corollary we conclude that  $\text{Aut}(\mathfrak{n})$  is maximal if and only if  $\mathfrak{n}$  is of Heisenberg type. These are complex Heisenberg algebras of various dimensions regarded as real Lie algebras.

First we recall the normal form for fat algebras with  $m = 2$  deduced from [LT]. Given  $c = a + bi \in \mathbb{C}$ , let

$$Z(c) = \begin{pmatrix} a & b \\ -b & a \end{pmatrix}.$$

If  $r \in \mathbb{Z}_+$ , set

$$A(c, r) = \begin{pmatrix} Z(c) & & & \\ I_2 & Z(c) & & \\ & & \ddots & \\ & & & I_2 & Z(c) \end{pmatrix}$$

a  $2r \times 2r$ -matrix. If  $\mathbf{c} = (c_1, \dots, c_\ell) \in \mathbb{C}^\ell$  and  $\mathbf{r} = (r_1, \dots, r_\ell) \in \mathbb{N}_+^\ell$ , set

$$A(\mathbf{c}, \mathbf{r}) = \begin{pmatrix} A(c_1, r_1) & & & \\ & A(c_2, r_2) & & \\ & & \ddots & \\ & & & A(c_\ell, r_\ell) \end{pmatrix}$$

which is a  $2s \times 2s$  matrix,  $s = r_1 + \dots + r_\ell$ .

Let now  $\phi, \psi_{(\mathbf{c}, \mathbf{r})}$  be the 2-forms on  $\mathbb{R}^{4s}$  whose matrices in the standard basis are

$$[\phi] = \begin{pmatrix} 0 & -I_{2s} \\ I_{2s} & 0 \end{pmatrix} \quad [\psi_{(\mathbf{c}, \mathbf{r})}] = \begin{pmatrix} 0 & A(\mathbf{c}, \mathbf{r}) \\ -A^\dagger(\mathbf{c}, \mathbf{r}) & 0 \end{pmatrix}. \tag{5}$$

Then

$$[u, v]_{(\mathbf{c}, \mathbf{r})} = (\phi(u, v), \psi_{(\mathbf{c}, \mathbf{r})}(u, v)) = \langle u, [\phi]v \rangle e_1 + \langle u, [\psi_{(\mathbf{c}, \mathbf{r})}]v \rangle e_2$$

is an  $\mathbb{R}^2$ -valued antisymmetric 2-form on  $\mathbb{R}^{4s}$ . Let

$$\mathfrak{n}_{(\mathbf{c}, \mathbf{r})} = \mathbb{R}^{4s} \oplus \mathbb{R}^2$$

be the corresponding Lie algebra.

Define  $M_{(\mathbf{c}, \mathbf{r})} \in \text{End}(\mathfrak{v})$  by

$$\phi(M_{(\mathbf{c}, \mathbf{r})}u, v) = \psi_{(\mathbf{c}, \mathbf{r})}(u, v),$$

whose matrix is

$$[M_{(\mathbf{c}, \mathbf{r})}] = \begin{pmatrix} -A^\dagger_{(\mathbf{c}, \mathbf{r})} & 0 \\ 0 & -A_{(\mathbf{c}, \mathbf{r})} \end{pmatrix}.$$

then we have

$$[u, v]_{(\mathbf{c}, \mathbf{r})} = \phi(u, v)e_1 + \phi(M_{(\mathbf{c}, \mathbf{r})}u, v)e_2, \text{ for } u, v \in \mathbb{R}^{4s}. \tag{6}$$

One can deduce [LT]

**Proposition 3.1.** (a) *Every fat algebra with center of dimension 2 is isomorphic to some  $\mathfrak{n}_{(\mathbf{c}, \mathbf{r})}$  with  $\mathbf{c} \in \mathbb{U}^\ell$ .*

(b) *Two of these are isomorphic if and only if the  $\mathbf{r}$ 's coincide up to permutations and the  $\mathbf{c}$ 's differ by some Möbius transformation acting componentwise.*

(c)  *$\mathfrak{n}_{(\mathbf{c}, \mathbf{r})}$  is of Heisenberg type if and only if  $\mathbf{c} = (c, \dots, c)$  and  $\mathbf{r} = (1, \dots, 1)$*

Let now

$$\mathfrak{n} = \mathfrak{n}_{(\mathbf{c}, \mathbf{r})}$$

be fat and let  $G = G(\mathfrak{n})$ , etc. We denote  $\hat{\mathfrak{n}}$  the algebra obtained by replacing the matrices  $A(c, r)$  by their semisimple parts and setting all  $c_j = \sqrt{-1}$ . The resulting

$\hat{A}(c, r)$  consists of blocks  $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$  along the diagonal and  $\hat{\mathfrak{n}}$  is isomorphic to the Heisenberg type algebra  $\mathfrak{n}_{((i, \dots, i), (1, \dots, 1))}$ . The correspondence

$$\mathfrak{n} \mapsto \hat{\mathfrak{n}}$$

is functorial and seems extendable inductively to fat algebras of any dimension, although here we will maintain the assumption  $m = 2$ .

**Lemma 3.2.**  $G_0(\mathfrak{n}) \subset G_0(\hat{\mathfrak{n}})$  and  $\dim \text{Aut}(\mathfrak{n}) \leq \dim \text{Aut}(\hat{\mathfrak{n}})$ .

**Proof.** Let  $\phi, \psi, M_{(\mathbf{c}, \mathbf{r})} \in \text{End}(\mathfrak{v})$  be as above, so that

$$\phi(M_{(\mathbf{c}, \mathbf{r})}u, v) = \psi_{(\mathbf{c}, \mathbf{r})}(u, v).$$

By formula (6),  $g \in G_0(\mathfrak{n}_{(\mathbf{c}, \mathbf{r})})$  if and only if

$$\phi(u, v) = \phi(gu, gv), \quad \phi(M_{(\mathbf{c}, \mathbf{r})}u, v) = \phi(M_{(\mathbf{c}, \mathbf{r})}gu, gv) = \phi(g^{-1}M_{(\mathbf{c}, \mathbf{r})}gu, v),$$

i.e., if and only if  $g \in \text{Sp}(\phi)$  and commutes with  $M_{(\mathbf{c}, \mathbf{r})}$ . In particular it commutes with the semisimple part  $\hat{M}_{(\mathbf{c}, \mathbf{r})}$ . This is conjugate to a matrix having blocks

$$Z(c) = \begin{pmatrix} \Re(c) & \Im(c) \\ -\Im(c) & \Re(c) \end{pmatrix} \text{ for various } c \in \mathbb{C} \text{ along the diagonal, and zeros elsewhere.}$$

Every matrix commuting with such a matrix will surely commute with that having all  $c = 1$ . It follows that  $g$  also preserves  $\phi(\hat{M}_{(\mathbf{c}, \mathbf{r})}u, v)$  and, therefore, it is an automorphism of  $\hat{\mathfrak{n}}$  as well. Thus,

$$G_0(\mathfrak{n}) \subset G_0(\hat{\mathfrak{n}}).$$

From Corollary 2.3,  $\dim(G(\mathfrak{n})/G_0(\mathfrak{n})) \leq \dim(G(\hat{\mathfrak{n}})/G_0(\hat{\mathfrak{n}}))$ , and therefore  $\dim G(\mathfrak{n}) = \dim(G(\mathfrak{n})/G_0(\mathfrak{n})) + \dim G_0(\mathfrak{n}) \leq \dim(G(\hat{\mathfrak{n}})/G_0(\hat{\mathfrak{n}})) + \dim G_0(\hat{\mathfrak{n}}) = \dim G(\hat{\mathfrak{n}})$ .

Formula (2) implies  $\dim \text{Aut}(\mathfrak{n}) \leq \dim \text{Aut}(\hat{\mathfrak{n}})$ , as claimed. ■

Next we will describe  $\mathfrak{g}_0(\mathfrak{n}_{(\mathbf{c}, \mathbf{r})})$  for  $c \in \mathbb{U}$  and  $r \in \mathbb{N}_+$ , i.e., the case when the matrices  $A$  consist of a single block. Since  $c$  is  $\text{SL}(2, \mathbb{C})$ -conjugate to  $i$ , it is enough to take  $c = i$ . Define the  $2 \times 2$ -matrices

$$\mathbf{1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \mathbf{i} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad \mathbf{x} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \mathbf{y} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix},$$

and let  $M_r(\mathbb{R}\langle \mathbf{1}, \mathbf{i} \rangle)$  and  $M_r(\mathbb{R}\langle \mathbf{x}, \mathbf{y} \rangle)$  denote the real vector spaces of  $r \times r$  matrices with coefficients in the span of  $\mathbf{1}, \mathbf{i}$  and  $\mathbf{x}, \mathbf{y}$  respectively. Then the vector space

$$\mathcal{R}(r) = \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} : A, D \in M_r(\mathbb{R}\langle \mathbf{1}, \mathbf{i} \rangle), B, C \in M_r(\mathbb{R}\langle \mathbf{x}, \mathbf{y} \rangle) \right\},$$

is actually a matrix algebra.

Note that

$$\mathbf{1}^t = \mathbf{1}, \quad \mathbf{i}^t = -\mathbf{i}, \quad \mathbf{x}^t = \mathbf{x}, \quad \mathbf{y}^t = \mathbf{y}.$$

Letting  $A^t$  denote the transpose of an  $\mathbb{R}$ -matrix and  $A^t, A^*$  the transpose and conjugate transpose of  $\mathbb{R}[\mathbf{i}, \mathbf{x}, \mathbf{y}]$ -matrices, one obtains

$$A^t = A^*$$

for  $A \in M_r(\mathbb{R}\langle \mathbf{1}, \mathbf{i} \rangle)$  while

$$A^t = A^t$$

for  $A \in M_r(\mathbb{R}\langle \mathbf{x}, \mathbf{y} \rangle)$ .

With the notation

$$J_1 = [\phi] \quad J_2 = [\psi_{((i,\dots,i),(1,\dots,1))}]$$

$$\mathfrak{g}_0(\hat{\mathfrak{n}}) = \{X \in \mathbb{R}^{4r \times 4r} : J_1 X + X^t J_1 = 0, J_2 X + X^t J_2 = 0\}.$$

From [S] we know that

$$\mathfrak{g}_0(\hat{\mathfrak{n}}) \cong \mathfrak{sp}(r, \mathbb{C})^{\mathbb{R}}$$

Changing basis,

$$\mathfrak{g}_0(\hat{\mathfrak{n}}) = \{X \in \mathcal{R}(r) : J_1 X + X^t J_1 = 0, J_2 X + X^t J_2 = 0\}$$

where

$$J_1 = \begin{pmatrix} 0 & I_r \\ -I_r & 0 \end{pmatrix}, \quad J_2 = \begin{pmatrix} 0 & \mathbf{i}I_r \\ \mathbf{i}I_r & 0 \end{pmatrix}.$$

This gives an alternative description of this algebra:

$$\mathfrak{g}_0(\hat{\mathfrak{n}}) = \left\{ \begin{pmatrix} A & B \\ C & -A^* \end{pmatrix} : A \in M_r(\mathbb{R}\langle \mathbf{1}, \mathbf{i} \rangle), B, C \in M_r(\mathbb{R}\langle \mathbf{x}, \mathbf{y} \rangle), B^t = B, C^t = C \right\}$$

We now restrict our attention to matrices  $\begin{pmatrix} A & B \\ C & -A^* \end{pmatrix}$  in  $\mathfrak{g}_0(\hat{\mathfrak{n}})$  where  $A, B, C$  have the respective forms

$$\begin{pmatrix} a_1 & a_2 & \cdots & a_r \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & a_2 \\ 0 & \cdots & 0 & a_1 \end{pmatrix} \quad \begin{pmatrix} b_1 & \cdots & b_{r-1} & b_r \\ \vdots & \ddots & \ddots & 0 \\ b_{r-1} & \ddots & \ddots & \vdots \\ b_r & 0 & \cdots & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & \cdots & 0 & c_1 \\ \vdots & \ddots & \ddots & c_2 \\ 0 & \ddots & \ddots & \vdots \\ c_1 & c_2 & \cdots & c_r \end{pmatrix}$$

with coefficients in  $\mathbb{R}^{2 \times 2}$ . Let  $\mathbf{A}_k = \begin{pmatrix} A & 0 \\ 0 & -A^* \end{pmatrix}$  having  $a_k = \mathbf{1}$  and zero otherwise and  $\mathbf{A}'_k$  the matrix of the same form but with  $a_k = \mathbf{i}$  and zeros elsewhere. Similarly, let  $\mathbf{B}_k$  (resp.  $\mathbf{C}_k$ ) the matrix  $\begin{pmatrix} 0 & B \\ 0 & 0 \end{pmatrix}$  (resp.,  $\begin{pmatrix} 0 & 0 \\ C & 0 \end{pmatrix}$ ) with  $b_k$  (resp.  $c_k$ ) equal to  $\mathbf{x}$  and zeros elsewhere, and  $\mathbf{B}'_k$  (resp.  $\mathbf{C}'_k$ ) with  $b_k$  (resp.  $c_k$ ) equal to  $\mathbf{y}$  and zeros elsewhere.

**Theorem 3.3.** *Let  $\mathfrak{n} = \mathfrak{n}_{(c,r)}$ ,  $(c,r) \in \mathbb{U} \times \mathbb{N}$ , and regard  $\mathfrak{g}_0(\mathfrak{n})$  as a subalgebra of  $\mathfrak{gl}(\mathfrak{v})$ . Then,*

1.  $\mathfrak{g}_0(\mathfrak{n})$  is the  $\mathbb{R}$ -span of  $\mathbf{A}_i, \mathbf{A}'_i, \mathbf{B}_i, \mathbf{B}'_i, \mathbf{C}_i, \mathbf{C}'_i$  for  $1 \leq i \leq r$ .
2. The semisimple part of  $\mathfrak{g}_0(\mathfrak{n})$  is the span of  $\mathbf{A}_1, \mathbf{A}'_1, \mathbf{B}_1, \mathbf{B}'_1, \mathbf{C}_1, \mathbf{C}'_1$ .
3. The solvable radical is the span of  $\mathbf{A}_i, \mathbf{A}'_i, \mathbf{B}_i, \mathbf{B}'_i, \mathbf{C}_i, \mathbf{C}'_i$  with  $1 < i \leq r$ .

In particular, the  $\mathbb{R}$ -dimension the  $\mathfrak{g}_0(\mathfrak{n})$  is equal to  $6r$  and the semisimple part of  $\mathfrak{g}_0(\mathfrak{n})$  is isomorphic to  $\mathfrak{sp}(1, \mathbb{C})$ .

**Proof.** It is enough to consider the case  $\mathfrak{n} = \mathfrak{n}_{(i,r)}$ . Let  $T_2 = [\psi_{(i,r)}]$  and write  $T_2 = J_2 + N_2$  where

$$N_2 = \begin{pmatrix} 0 & N \\ -N^t & 0 \end{pmatrix}, \text{ with } N = \begin{pmatrix} 0 & \cdots & 0 & 0 \\ 1 & \ddots & 0 & 0 \\ & \ddots & \ddots & \\ 0 & \cdots & 1 & 0 \end{pmatrix}.$$

From Lemma 3.2,  $\mathfrak{g}_0(\mathfrak{n}) = \{X \in \mathfrak{g}_0(\hat{\mathfrak{n}}) : T_2X + X^tT_2 = 0\}$ . As  $\mathfrak{g}_0(\mathfrak{n}) \subset \mathfrak{g}_0(\hat{\mathfrak{n}})$  one obtains

$$\mathfrak{g}_0(\mathfrak{n}) = \{X \in \mathfrak{g}_0(\hat{\mathfrak{n}}) : N_2X + X^tN_2 = 0\}.$$

The conditions on  $\begin{pmatrix} A & B \\ C & -A^* \end{pmatrix} \in \mathfrak{g}_0(\mathfrak{n})$  are, explicitly,

$$0 = NC - C^tN^t = NC - (NC)^t \tag{7}$$

$$0 = N^tA - AN^t \tag{8}$$

$$0 = N^tB - B^tN = N^tB - (N^tB)^t. \tag{9}$$

For the first equation, note that  $NC$  symmetric if and only if  $c_{i,j+1} = c_{j,i+1}$  and  $c_{1,j} = 0$  for  $i, j < n$ . Since  $C$  is symmetric,  $c_{i,j+1} = c_{j,i+1} = c_{i+1,j}$  and  $c_{1,j} = 0$  for  $i, j < n$ . We conclude:

$$\text{If } i + j = k \leq r, c_{i,j} = c_{i,k-i} = c_{i-1,k-i+1} = c_{i-2,k-i+2} \cdots = c_{1,k-1} = 0$$

$$\text{If } i + j = k > r, c_{i,j} = c_{i,k-i} = c_{i+1,k-i-1} = c_{i+2,k-i-2} \cdots = c_{r,k-i+i-r} = c_{r,k-r}$$

Thus, the strict upper antidiagonals are zero and each lower antidiagonal have all its elements equal.

For the second equation, note that  $N^t$  and  $A$  commute. This is equivalent to  $c_{i,j} = c_{t,s}$  when  $j - i = s - t$  and  $c_{i,1} = 0$  for  $i > 1$ . The first condition implies that each diagonal have all its elements equal, while the second implies that the strict lower diagonals are zero.

Equation (9) is analogous to equation (7): the condition  $N^tB$  symmetric is equivalent to each antidiagonal have all its elements equal and that the strict lower antidiagonals are 0.

From all this we conclude that the span of  $\mathbf{A}_i, \mathbf{A}'_i, \mathbf{B}_i, \mathbf{B}'_i, \mathbf{C}_i, \mathbf{C}'_i$  with  $1 \leq i \leq r$  is  $\mathfrak{g}_0(\mathfrak{n})$  and (1) follows.

(2) and (3) follow from (1) and the explicit presentation of the matrices  $\mathbf{A}_i, \mathbf{A}'_i, \mathbf{B}_i, \mathbf{B}'_i, \mathbf{C}_i, \mathbf{C}'_i$ . ■

**Corollary 3.4.** (of the proof) Let  $\mathfrak{n}$  be fat. Then  $\dim(\mathfrak{g}_0(\mathfrak{n}))$  is maximal if and only if  $\mathfrak{n}$  is of Heisenberg type.

**Proof.** Let  $(\mathbf{c}, \mathbf{r}) = ((c_1, \dots, c_l), (r_1, \dots, r_l))$  be such that  $\mathfrak{n} = \mathfrak{n}_{(\mathbf{c}, \mathbf{r})}$ . We know that  $\mathfrak{g}_0(\mathfrak{n}) \subset \mathfrak{g}_0(\hat{\mathfrak{n}})$ . If  $c_i \neq c_j$  for some  $i, j$ , then there is not intertwining operator between the blocks corresponding to these invariants, so  $\mathfrak{g}_0(\mathfrak{n}) \neq \mathfrak{g}_0(\hat{\mathfrak{n}})$ .

When  $c_1 = c_2 = \dots = c_\ell$  we can consider  $c_j = i$  for all  $j$ . Let  $r = \sum r_i$ . In this case if  $\begin{pmatrix} A & B \\ C & -A^* \end{pmatrix} \in \mathfrak{g}_0(\mathbf{n})$  must satisfy the equations (7), (8), (9) but with  $N$  such that coefficients  $n_{j+1,j}$  are 0 or  $\mathbf{1}$ . Suppose now that  $\mathfrak{g}_0(\mathbf{n})$  is not of Heisenberg type, then some  $n_{j+1,j}$  is equal to  $\mathbf{1}$ . We assume that  $n_{21} = \mathbf{1}$  and let  $A \in M_r(\mathbb{R}(\mathbf{1}, \mathbf{i}))$  such that  $a_{12} = \mathbf{1}$  and 0 otherwise, then

$$X = \begin{pmatrix} A & 0 \\ 0 & -A^* \end{pmatrix}$$

belongs to  $\mathfrak{g}_0(\hat{\mathbf{n}})$  but is not in  $\mathfrak{g}_0(\mathbf{n})$ . ■

It can be shown in general that the semisimple part of  $\mathfrak{g}_0(\mathbf{n})$  is isomorphic to  $\oplus_i \mathfrak{sp}(m_i, \mathbb{C})$ , where  $m_i$  is the multiplicity of the pair  $(c_i, r_i)$  in  $(\mathbf{c}, \mathbf{r})$ .

In the case  $m = 2$ ,  $\mathfrak{g}/\mathfrak{g}_0$  is either 0 or isomorphic to  $\mathfrak{so}(2)$ .

**Theorem 3.5.**  $\mathfrak{g}(\mathbf{n})/\mathfrak{g}_0(\mathbf{n}) \cong \mathfrak{so}(2)$  if  $c_1 = \dots = c_\ell$ , and 0 otherwise.

**Proof.**  $\mathfrak{g}/\mathfrak{g}_0$  is a compact subalgebra of  $\mathfrak{gl}(2)$ , hence of the form  $g\mathfrak{so}(2)g^{-1}$  for some  $g \in \text{SL}(2, \mathbb{R})$  and it is nonzero if and only if there exists  $X \in \mathfrak{sl}(v)$  such that, in the notation of the proof of Theorem 3.3,

$$\begin{pmatrix} X & 0 \\ 0 & g\mathbf{i}g^{-1} \end{pmatrix}$$

is a derivation of  $\mathbf{n}$ . For  $g = \mathbf{1}$ , if  $T_1, T_2$  correspond to the standard basis of  $\mathfrak{z}$ , the equations for  $X$  become

$$(a) \quad T_1X + X^tT_1 = T_2, \quad (b) \quad T_2X + X^tT_2 = -T_1$$

In normal form, and for a single block  $A_{(i,r)}$ ,

$$T_1 = J_1 = \begin{pmatrix} 0 & I_r \\ -I_r & 0 \end{pmatrix}, \quad T_2 = \begin{pmatrix} 0 & \mathbf{i}I_r + N \\ \mathbf{i}I_r - N^t & 0 \end{pmatrix}.$$

We decompose

$$T_2 = J_2 + N_2, \quad \text{with} \quad J_2 = \begin{pmatrix} 0 & \mathbf{i}I_r \\ \mathbf{i}I_r & 0 \end{pmatrix}, \quad N_2 = \begin{pmatrix} 0 & N \\ -N^t & 0 \end{pmatrix}$$

and regard  $J_1, J_2, T_1, T_2, N_2$  as matrices with coefficients in  $\mathbb{R}^{2 \times 2}$ . Note that  $J_1, J_2$  correspond to  $\hat{\mathbf{n}}$ , of Heisenberg type. Let

$$Y_0 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & s & 0 \\ 0 & 2\mathbf{i} & 0 & 0 & 0 & & s & 0 \\ 0 & \mathbf{1} & 4\mathbf{i} & 0 & 0 & 0 & s & 0 \\ 0 & 0 & 2\mathbf{1} & 6\mathbf{i} & 0 & 0 & s & 0 \\ 0 & 0 & 0 & 3\mathbf{1} & 8\mathbf{i} & & s & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & 0 \\ 0 & 0 & 0 & \vdots & \vdots & 0 & (n-2)\mathbf{1} & 2(n-1)\mathbf{i} \end{pmatrix}.$$

A straightforward calculation shows that

$$X_0 = \begin{pmatrix} -Y_0^t & 0 \\ 0 & -Y_0^t + \mathbf{i}I_r + N \end{pmatrix}$$

is a solution of (a), (b). We conclude that

$$\begin{pmatrix} X_0 & 0 \\ 0 & \mathbf{i} \end{pmatrix}$$

is a derivation of  $\mathfrak{n}_{(i,r)}$ , which lies in  $\mathfrak{g}(\mathfrak{n}_{(i,r)})$  but not in  $\mathfrak{g}_0(\mathfrak{n}_{(i,r)})$ .

For any  $c \in \mathbb{U}$ ,  $\mathfrak{n}_{(c,r)} \cong \mathfrak{n}_{(i,r)}$ , hence they have the same  $\mathfrak{g}/\mathfrak{g}_0$  up to isomorphisms. In fact, for any  $g \in \text{SL}(2, \mathbb{R})$ , the algebra  $\mathfrak{n}_{(g \cdot i, r)}$  has a derivation of the form

$$\begin{pmatrix} X & 0 \\ 0 & g\mathbf{i}g^{-1} \end{pmatrix}.$$

For a fixed  $g$ , these  $X$  are unique modulo  $\mathfrak{g}_0$  and come in normal form. Clearly,  $c$  determines the  $2 \times 2$  matrix  $g\mathbf{i}g^{-1}$  and the complex number  $g \cdot i$ .

In the case of an arbitrary fat  $\mathfrak{n}_{(c,r)}$ , each block  $(c_k, r_k)$  determines a corresponding  $X_k$  such that

$$\begin{pmatrix} X_k & 0 \\ 0 & g_k\mathbf{i}g_k^{-1} \end{pmatrix}$$

is a derivation of  $\mathfrak{n}_{(c_k, r_k)}$ . If  $\mathfrak{n}_{(c,r)}$  has a derivation in  $\mathfrak{g}$  that is not in  $\mathfrak{g}_0$ , then its must have one which is combination of such, acting on  $\mathfrak{v}$  as  $X_1 + X_2 + \dots$ . This forces all the  $g_k\mathbf{i}g_k^{-1}$  to be the same and all the  $c_i$  to be the same. The reciprocal is clear. ■

In particular, all algebras  $\mathfrak{n}_{(c,r)}$  with  $c_1 = \dots = c_\ell$  and  $r_i > 1$  maximize the dimension of  $\mathfrak{g}/\mathfrak{g}_0$ , but they are not Heisenberg type.

Lauret had pointed out to us that there were non Heisenberg type algebras such that  $\mathfrak{g}(\mathfrak{n})/\mathfrak{g}_0(\mathfrak{n}) \neq 0$ . Independently, Oscari proved that this holds whenever the  $c_i$ 's all agree.

#### 4. Fat distributions

Let  $D$  be a smooth vector distribution on a smooth manifold  $M$ , i.e., a subbundle of the tangent bundle  $T(M)$ . Its nilpotentization, or symbol, is the bundle on  $M$  with fiber

$$N^D(M)_p = \bigoplus_j D_p^{(j)} / D_p^{(j-1)}$$

where  $D_p^{(1)} = D_p$  and  $D_p^{(j+1)} = D_p^{(j)} + [\Gamma(D), \Gamma(D^j)]_p$ . The Lie bracket in  $\Gamma(T(M))$  induces a graded nilpotent Lie algebra structure on each fiber of  $N^D(M)$ . If  $D^{(j)} = T(M)$  for some  $j$ ,  $D$  is called completely non-integrable. If  $D^{(2)} = T(M)$ , the nilpotentization is 2-step, which in the notation of the previous section, is

$$\mathfrak{n}_p = N_D(M)_p = D_p \oplus \frac{D_p + [\Gamma(D), \Gamma(D)]_p}{D_p} = \mathfrak{v}_p + \mathfrak{z}_p,$$

It is also easy to see that  $D$  is fat in the sense of Weinstein [Mo] if and only if  $\mathfrak{n}_p = \mathfrak{v} + \mathfrak{z}$  is non-singular, i.e., fat in the sense defined in the section 1.

A subriemannian metric  $g$  defined on  $D$  determines a metric on  $\mathfrak{v}$ . On  $\mathfrak{z}$  we put a metric  $\sigma$  invariant under  $G$ . Let  $\{\phi_1, \dots, \phi_m; \psi_1, \dots, \psi_n\}$  be a coframe on  $M$  such that

$$D = \cap \ker \phi_i,$$

with  $\{\phi_1, \dots, \phi_m\}$  and  $\{\psi_1, \dots, \psi_n\}$  orthonormal with respect to  $g + \sigma$ . Define  $T_z \in \text{End}(D)$  as before, by

$$\sigma(z, [u, v]) = g(T_z u, v).$$

Then  $D$  is fat if and only if  $T_z$  is invertible for all non-zero  $z \in \mathfrak{z}$ . The structure equations for the coframe can be written

$$d\phi_k \equiv \sum_i (T_k \psi_i) \wedge \psi_i \quad \text{mod}(\phi_\ell)$$

with the  $T_k$ 's having the property that any non-zero linear combination of them is invertible. This is deduced from the fact that if  $u, v \in \mathfrak{v}$ , then  $d\phi[u, v] = -\phi([u, v])$ , since  $u(\phi(v)) = u(0) = 0$ . The  $d\psi$ 's are essentially arbitrary.

Let now  $M$  be a the simply connected Lie group with a fat Lie algebra  $\mathfrak{n}$ ,  $D$  the left-invariant distribution on  $M$  such that  $D_e = \mathfrak{v}$ . For a left-invariant coframe, the structure equations take the form

$$d\phi_k = \sum_i (J_k \psi_i) \wedge \psi_i, \quad d\psi_i = 0$$

where  $J_1, \dots, J_m$  are anticommuting complex structures on  $D$ .

The results from the previous sections lead to consider fat distributions satisfying

$$(4.1) \quad d\phi_k = \sum_i (J_k \psi_i) \wedge \psi_i \quad \text{mod}(\phi_\ell)$$

where the  $J_k$  are sections of  $\text{End}(T(M)^*)$  satisfying the Canonical Commutation Relations

$$J_i J_j + J_j J_i = -2\delta_{ij}.$$

The Equivalence Problem for these systems has been discussed for distributions with growth vector  $(2n, 2n + 1), (4n, 4n + 3)$  and  $(8, 15)$ . In these cases  $\mathfrak{n}$  is parabolic, i.e., isomorphic to the Iwasawa subalgebra of a real semisimple Lie algebra  $\mathfrak{g}$  of real rank one. The Tanaka [T] subriemannian prolongation of such algebra is  $\mathfrak{g}$ , while in the non-parabolic case is just

$$\mathfrak{n} + \mathfrak{k}(\mathfrak{n}) + \mathfrak{a}(\mathfrak{n})$$

where  $\mathfrak{a}(\mathfrak{n})$  the 1-dimensional Lie algebra of dilations [Su]. In this case, Tanaka's theorem implies that, in the notation of [Z], the first pseudo G-structure  $P^0$  already carries a canonical frame.

As this paper was being written, E. van Erp pointed out to us his article [Er], where fat distributions are called polycontact and those satisfying (4.1) arise by imposing a compatible conformal structure.

**Acknowledgments.** We wish to thank Professor J. Vargas for helpful discussions and M. Subils for pointing out a mistake in a previous version of this paper.

### References

- [BTV] Berndt, J., F. Tricerri and L. Vanhecke, “Generalized Hesienberg groups and Damek-Ricci Harmonic spaces,” Springer Lecture Notes **1598**, Berlin, 1995.
- [CGN] Capogna, L., N. Garofalo, and D. M. Nhieu, *Properties of harmonic measures in the Dirichlet Problem for Lie groups of Heisenberg type*, Amer. J. Math **124** (2002), 273–306.
- [DG] de Graaf, W., “Lie algebras: theory and algorithms,” North Holland Math. Library **56** Elsevier, 2000.
- [E] Eberlein, P., *Geometry of 2-step nilpotent groups with a left-invariant metric*, Annales Scientifiques de l’E.N.S. (4)**27** (1994), 611–660.
- [Er] van Erp, E., *Contact structures of arbitrary codimension and idempotents in the Heisenberg algebra*, arXiv:1001.5426v1 [math.DG] (2010)
- [GV] Garofalo, N., and D. Vassiliev, *Symmetry properties of solutions of Yamabe’s equation on groups of Heisenberg type*, Duke Math. J. **106** (2001), 411–448.
- [H] Husemoller, D., “Fibre Bundles,” Graduate texts in mathematics **20**, Springer-Verlag, New York, 1994.
- [K] Kaplan, A., *Fundamental solutions for a class of hypoelliptic PDE generated by composition of quadratic forms*, Trans. Amer. Math. Soc. **258** (1980), 147–153.
- [LT] Levstein, F., and A. Tiraboschi, *Classes of 2-step nilpotent Lie algebras*, Comm. in Algebra **27** (1999), 2425–2440.
- [M] Métivier, G., *Hypoellipticite analytique sur des groupes nilpotents de rang 2*, Duke Math. J. **47** (1980), 195–221.
- [Mo] Montgomery, R., *A tour of subriemannian geometries, their geodesics and applications*, Amer. Math. Soc. Mathematical Surveys and Monographs , Providence, R. I., 2002.
- [MS] Müller, D. and A. Seeger, *Singular spherical maximal operators on a class of two step nilpotent Lie groups*, Israel J. Math. **141** (2004), 315–390.

- [P] Pansu, P., *Métriques de Carnot-Carathodory et quasiisométries des espaces symétriques de rang un*, Ann. of Math. **129** (1989), 1–60.
- [S] Saal, L., *The automorphism group of a Lie algebra of Heisenberg type*, Rend. Sem. Mat. Univ. Pol. Torino **54**(1996), 101–113.
- [Su] Subils, M., Personal communication.
- [T] Tanaka, N., *On differential systems, graded Lie algebras and pseudogroups* J. Math. Kyoto. Univ. **10** (1970), 1–82.
- [Z] Zelenko, I., *On Tanaka's prolongation procedure for filtered structures of constant type* SIGMA **5** (2009), Paper 094, 21pp.

Aroldo Kaplan  
Facultad de Matemática,  
Astronomía y Física  
Universidad Nacional de Córdoba  
CIEM – CONICET. (5000)  
Ciudad Universitaria  
Córdoba, Argentina  
kaplan@famaf.unc.edu.ar

Alejandro Tiraboschi  
Facultad de Matemática,  
Astronomía y Física  
Universidad Nacional de Córdoba  
CIEM – CONICET. (5000)  
Ciudad Universitaria  
Córdoba, Argentina  
tirabo@famaf.unc.edu.ar

Received November 22, 2011  
and in final form March 15, 2013