

Kac-Moody Lie Algebras Graded by Kac-Moody Root Systems

Hechmi Ben Messaoud and Guy Rousseau

Communicated by A. Valette

Abstract. We look to gradations of Kac-Moody Lie algebras by Kac-Moody root systems with finite dimensional weight spaces. We extend, to general Kac-Moody Lie algebras, the notion of C -admissible pair as introduced by H. Rubenthaler and J. Nervi for semi-simple and affine Lie algebras. If \mathfrak{g} is a Kac-Moody Lie algebra (with Dynkin diagram indexed by I) and (I, J) is such a C -admissible pair, we construct a C -admissible subalgebra \mathfrak{g}^J , which is a Kac-Moody Lie algebra of the same type as \mathfrak{g} , and whose root system Σ grades finitely the Lie algebra \mathfrak{g} . For an admissible quotient $\rho : I \rightarrow \bar{I}$ we build also a Kac-Moody subalgebra \mathfrak{g}^ρ which grades finitely the Lie algebra \mathfrak{g} . If \mathfrak{g} is affine or hyperbolic, we prove that the classification of the gradations of \mathfrak{g} is equivalent to those of the C -admissible pairs and of the admissible quotients. For general Kac-Moody Lie algebras of indefinite type, the situation may be more complicated; it is (less precisely) described by the concept of generalized C -admissible pairs.

Mathematics Subject Classification 2000: 17B67.

Key Words and Phrases: Kac-Moody algebra, C -admissible pair, gradation.

Introduction

The notion of gradation of a Lie algebra \mathfrak{g} by a finite root system Σ was introduced by S. Berman and R. Moody [8] and further studied by G. Benkart and E. Zelmanov [5], E. Neher [15], B. Allison, G. Benkart and Y. Gao [1] and J. Nervi [16]. This notion was extended by J. Nervi [17] to the case where \mathfrak{g} is an affine Kac-Moody algebra and Σ the (infinite) root system of an affine Kac-Moody algebra; in her two articles she uses the notion of C -admissible subalgebra associated to a C -admissible pair for the Dynkin diagram, as introduced by H. Rubenthaler [21].

We consider here a general Kac-Moody algebra \mathfrak{g} (indecomposable and symmetrizable) and the root system Σ of a Kac-Moody algebra. We say that \mathfrak{g} is finitely Σ -graded if \mathfrak{g} contains a Kac-Moody subalgebra \mathfrak{m} (the grading subalgebra) whose root system relatively to a Cartan subalgebra \mathfrak{a} of \mathfrak{m} is Σ and moreover the action of $ad(\mathfrak{a})$ on \mathfrak{g} is diagonalizable with weights in $\Sigma \cup \{0\}$ and

finite dimensional weight spaces, see Definition 1.4. The finite dimensionality of weight spaces is a new condition, it was fulfilled by the non-trivial examples of J. Nervi [17] but it excludes the gradings of infinite dimensional Kac-Moody algebras by finite root systems as in [5]. Many examples of these gradations are provided by the almost split real forms of \mathfrak{g} , cf. 1.6. We are interested in describing the possible gradations of a given Kac-Moody algebra (as in [16], [17]), not in determining all the Lie algebras graded by a given root system Σ (as e.g. in [1] for Σ finite). We carry out completely this project when \mathfrak{g} is affine or hyperbolic.

Let I be the index set of the Dynkin diagram of \mathfrak{g} , we generalize the notion of C -admissible pair (I, J) as introduced by H. Rubenthaler [21] and J. Nervi [16], [17], cf. Definition 2.1. For each Dynkin diagram I the classification of the C -admissible pairs (I, J) is easy to deduce from the list of irreducible C -admissible pairs due to these authors. We are able then to generalize in section 2 their construction of a C -admissible subalgebra (associated to a C -admissible pair) which grades finitely \mathfrak{g} :

Theorem 1. (cf. 2.6, 2.11, 2.14) *Let \mathfrak{g} be an indecomposable and symmetrizable Kac-Moody algebra, associated to a generalized Cartan matrix $A = (a_{i,j})_{i,j \in I}$. Let $J \subset I$ be a subset of finite type such that the pair (I, J) is C -admissible. There is a generalized Cartan matrix $A^J = (a'_{k,l})_{k,l \in I'}$ with index set $I' = I \setminus J$ and a Kac-Moody subalgebra \mathfrak{g}^J of \mathfrak{g} associated to A^J , with root system Δ^J . Then \mathfrak{g} is finitely Δ^J -graded with grading subalgebra \mathfrak{g}^J .*

For a general finite gradation of \mathfrak{g} with grading subalgebra \mathfrak{m} , we prove (in section 3) that \mathfrak{m} also is indecomposable, symmetrizable and the restriction to \mathfrak{m} of the invariant bilinear form of \mathfrak{g} is non-degenerate (3.11 and 3.17). The Kac-Moody algebras \mathfrak{g} and \mathfrak{m} have the same type: finite, affine or indefinite; the first two types correspond to the cases already studied e.g. by J. Nervi. Moreover if \mathfrak{g} is indefinite Lorentzian or hyperbolic, then so is \mathfrak{m} (Propositions 3.6 and 3.27). We get also the following precise structure result for this general situation :

Theorem 2. *Let \mathfrak{g} be an indecomposable and symmetrizable Kac-Moody algebra, finitely graded by a root system Σ of Kac-Moody type with grading subalgebra \mathfrak{m} .*

1) *We may choose the Cartan subalgebras \mathfrak{a} of \mathfrak{m} , \mathfrak{h} of \mathfrak{g} such that $\mathfrak{a} \subset \mathfrak{h}$. Then there is a surjective map $\rho_a : \Delta \cup \{0\} \rightarrow \Sigma \cup \{0\}$ between the corresponding root systems. We may choose the bases $\Pi_a = \{\gamma_s \mid s \in \bar{I}\} \subset \Sigma$ and $\Pi = \{\alpha_i \mid i \in I\} \subset \Delta$ of these root systems such that $\rho_a(\Delta^+) \subset \Sigma^+ \cup \{0\}$ and $\{\alpha \in \Delta \mid \rho_a(\alpha) = 0\} = \Delta_J := \Delta \cap (\sum_{j \in J} \mathbb{Z}\alpha_j)$ for some subset $J \subset I$ of finite type.*

2) *Let $I'_{re} = \{i \in I \mid \rho_a(\alpha_i) \in \Pi_a\}$, $I'_{im} = \{i \in I \mid \rho_a(\alpha_i) \notin \Pi_a \cup \{0\}\}$. Then $J = \{i \in I \mid \rho_a(\alpha_i) = 0\}$. We note I_{re} (resp. J°) the union of the connected components of $I \setminus I'_{im} = I'_{re} \cup J$ meeting I'_{re} (resp. contained in J), and $J_{re} = J \cap I_{re}$. Then the pair (I_{re}, J_{re}) is C -admissible (eventually decomposable).*

3) *There is a Kac-Moody subalgebra $\mathfrak{g}(I_{re})$ of \mathfrak{g} , associated to I_{re} , which contains \mathfrak{m} . This Kac-Moody Lie algebra is finitely $\Delta(I_{re})^{J_{re}}$ -graded, with grading subalgebra $\mathfrak{g}(I_{re})^{J_{re}}$. Both algebras $\mathfrak{g}(I_{re})$ and $\mathfrak{g}(I_{re})^{J_{re}}$ are finitely Σ -graded with grading subalgebra \mathfrak{m} .*

It may happen that I'_{im} is non-empty, we then say that (I, J) is a gener-

alized C -admissible pair and the gradation is imaginary. We give and explain precisely an example in section 5.

When I'_{im} is empty (i.e. when the gradation is real : 3.16), $I_{re} = I$, $J_{re} = J$, $\mathfrak{g}(I_{re}) = \mathfrak{g}$, $(I, J) = (I_{re}, J_{re})$ is a C -admissible pair and the situation looks much like the one described by J. Nervi in the finite [16] or affine [17] cases. Actually we prove that this is always true when \mathfrak{g} is of finite type, affine or hyperbolic (Proposition 3.26). In this real case we get the gradation of \mathfrak{g} with two levels: \mathfrak{g} is finitely Δ^J -graded with grading subalgebra \mathfrak{g}^J as in Theorem 1 and \mathfrak{g}^J is finitely Σ -graded with grading subalgebra \mathfrak{m} . But the gradation of \mathfrak{g}^J by Σ and \mathfrak{m} is such that the corresponding set " J " described as in Theorem 2 is empty; we say (following [16], [17]) that it is a maximal gradation, cf. Definition 3.16 and Proposition 3.21.

To get a complete description of the real gradations, it remains to describe the maximal gradations; this is done in section 4. We prove in Proposition 4.1 that a maximal gradation $(\mathfrak{g}, \Sigma, \mathfrak{m})$ is entirely described by a quotient map $\rho : I \rightarrow \bar{I}$ which is admissible i.e. satisfies two simple conditions (MG1) and (MG2) with respect to the generalized Cartan matrix $A = (a_{i,j})_{i,j \in I}$. Conversely for any admissible quotient map ρ , it is possible to build a maximal gradation of \mathfrak{g} associated to this map, cf. Proposition 4.5 and Remark 4.7.

1. Preliminaries

We recall the basic results on the structure of Kac-Moody Lie algebras and we set the notations. More details can be found in the book of Kac [12]. We end by the definition of finitely graded Kac-Moody algebras.

Generalized Cartan matrices

Let I be a finite index set. A matrix $A = (a_{i,j})_{i,j \in I}$ is called a *generalized Cartan matrix* if it satisfies :

- (1) $a_{i,i} = 2 \quad (i \in I)$
- (2) $a_{i,j} \in \mathbb{Z}^- \quad (i \neq j)$
- (3) $a_{i,j} = 0$ implies $a_{j,i} = 0$.

The matrix A is called *decomposable* if for a suitable permutation of I it takes the form $\begin{pmatrix} B & 0 \\ 0 & C \end{pmatrix}$ where B and C are square matrices. If A is not decomposable, it is called *indecomposable*.

The matrix A is called *symmetrizable* if there exists an invertible diagonal matrix $D = \text{diag}(d_i, i \in I)$ such that DA is symmetric. The entries $d_i, i \in I$, can be chosen to be positive rationals and if moreover the matrix A is indecomposable, then these entries are unique up to a constant factor.

Any indecomposable generalized Cartan matrix is of one of three mutually exclusive types : *finite*, *affine* and *indefinite* ([12, Chap. 4]). A generalized Cartan matrix is said of *finite type* if each of its indecomposable factors is of finite type. An indecomposable and symmetrizable generalized Cartan matrix A is called *Lorentzian* if it is non-singular and the corresponding symmetric matrix has signature $(+ + \dots + -)$; it is then of indefinite type.

An indecomposable generalized Cartan matrix A is called *strictly hyperbolic* (resp. *hyperbolic*) if the deletion of any one vertex, and the edges connected to it, of the corresponding Dynkin diagram yields a disjoint union of Dynkin diagrams of finite (resp. finite or affine) type.

Note that a symmetrizable hyperbolic generalized Cartan matrix is non-singular and Lorentzian (cf. [14]).

Kac-Moody algebras and groups (See [12] and [18]).

Let $A = (a_{i,j})_{i,j \in I}$ be a symmetrizable generalized Cartan matrix. Let $(\mathfrak{h}_{\mathbb{R}}, \Pi = \{\alpha_i, i \in I\}, \Pi^{\check{}} = \{\alpha_i^{\check{}}, i \in I\})$ be a realization of A over the real field \mathbb{R} : thus $\mathfrak{h}_{\mathbb{R}}$ is a real vector space such that $\dim(\mathfrak{h}_{\mathbb{R}}) = |I| + \text{corank}(A)$, Π and $\Pi^{\check{}}$ are linearly independent in $\mathfrak{h}_{\mathbb{R}}^*$ and $\mathfrak{h}_{\mathbb{R}}$ respectively such that $\langle \alpha_j, \alpha_i^{\check{}} \rangle = a_{i,j}$. Let $\mathfrak{h} = \mathfrak{h}_{\mathbb{R}} \otimes \mathbb{C}$, then $(\mathfrak{h}, \Pi, \Pi^{\check{}})$ is a realization of A over the complex field \mathbb{C} . It follows that, if A is non-singular, then $\Pi^{\check{}}$ (resp. Π) is a basis of \mathfrak{h} (resp. \mathfrak{h}^*); moreover $\mathfrak{h}_{\mathbb{R}} = \{h \in \mathfrak{h} \mid \alpha_i(h) \in \mathbb{R}, \forall i \in I\}$ is well defined by the realization $(\mathfrak{h}, \Pi, \Pi^{\check{}})$.

Let $\mathfrak{g} = \mathfrak{g}(A)$ be the complex Kac-Moody Lie algebra associated to A : it is generated by $\{\mathfrak{h}, e_i, f_i, i \in I\}$ with the following relations

$$\begin{aligned} [\mathfrak{h}, \mathfrak{h}] &= 0, & [e_i, f_j] &= \delta_{i,j} \alpha_i^{\check{}} & (i, j \in I); \\ [h, e_i] &= \langle \alpha_i, h \rangle e_i, & [h, f_i] &= -\langle \alpha_i, h \rangle f_i & (h \in \mathfrak{h}); \\ (\text{ad } e_i)^{1-a_{i,j}}(e_j) &= 0, & (\text{ad } f_i)^{1-a_{i,j}}(f_j) &= 0 & (i \neq j). \end{aligned} \tag{1.1}$$

The Kac-Moody algebra $\mathfrak{g} = \mathfrak{g}(A)$ decomposes as a direct sum of factors $\mathfrak{g}(A_i)$, where A_1, \dots, A_r are the indecomposable factors of A . It is said indecomposable if the corresponding generalized Cartan matrix A is indecomposable and of finite, affine or indefinite type if A is.

The derived algebra \mathfrak{g}' of \mathfrak{g} is generated by the *Chevalley generators* $e_i, f_i, i \in I$, and the center \mathfrak{c} of \mathfrak{g} lies in $\mathfrak{h}' = \mathfrak{h} \cap \mathfrak{g}' = \sum_{i \in I} \mathbb{C} \alpha_i^{\check{}}$. If the generalized Cartan matrix A is indecomposable and non-singular, then $\mathfrak{g} = \mathfrak{g}'$ is a (finite or infinite)-dimensional simple Lie algebra, and the center \mathfrak{c} is trivial.

The subalgebra \mathfrak{h} is a maximal $\text{ad}(\mathfrak{g})$ -diagonalizable subalgebra of \mathfrak{g} , it is called the *standard Cartan subalgebra* of \mathfrak{g} . Let $\Delta = \Delta(\mathfrak{g}, \mathfrak{h})$ be the corresponding root system; then Π is a root basis of Δ and $\Delta = \Delta^+ \cup \Delta^-$, where $\Delta^{\pm} = \Delta \cap \mathbb{Z}^{\pm} \Pi$ is the set of positive (or negative) roots relative to the basis Π . For $\alpha \in \Delta$, let \mathfrak{g}_{α} be the root space of \mathfrak{g} corresponding to the root α ; then $\mathfrak{g} = \mathfrak{h} \oplus (\bigoplus_{\alpha \in \Delta} \mathfrak{g}_{\alpha})$.

The *Weyl group* W of $(\mathfrak{g}, \mathfrak{h})$ is generated by the fundamental reflections r_i ($i \in I$) such that $r_i(h) = h - \langle \alpha_i, h \rangle \alpha_i^{\check{}}$ for $h \in \mathfrak{h}$, it is a Coxeter group on $\{r_i, i \in I\}$ with length function $w \mapsto l(w), w \in W$. The Weyl group W acts on \mathfrak{h}^* and Δ , we set $\Delta^{re} = W(\Pi)$ (the real roots) and $\Delta^{im} = \Delta \setminus \Delta^{re}$ (the imaginary roots). If the generalized Cartan matrix A is indecomposable, then any root basis of Δ is W -conjugate to Π or $-\Pi$.

A *Borel subalgebra* of \mathfrak{g} is a maximal completely solvable subalgebra. A *parabolic subalgebra* of \mathfrak{g} is a (proper) subalgebra containing a Borel subalgebra. The *standard positive (or negative) Borel subalgebra* is $\mathfrak{b}^{\pm} := \mathfrak{h} \oplus (\bigoplus_{\alpha \in \Delta^{\pm}} \mathfrak{g}_{\alpha})$. A parabolic subalgebra \mathfrak{p}^+ (resp. \mathfrak{p}^-) containing \mathfrak{b}^+ (resp. \mathfrak{b}^-) is called *positive (resp. negative) standard parabolic subalgebra* of \mathfrak{g} ; then there exists a subset J

of I (called the type of \mathfrak{p}^\pm) such that $\mathfrak{p}^\pm = \mathfrak{p}^\pm(J) := (\bigoplus_{\alpha \in \Delta_J} \mathfrak{g}_\alpha) + \mathfrak{b}^\pm$, where $\Delta_J = \Delta \cap (\bigoplus_{j \in J} \mathbb{Z}\alpha_j)$ (cf. [13]).

In [18], D.H. Peterson and V.G. Kac construct a group G , which is the connected and simply connected complex algebraic group associated to \mathfrak{g} when \mathfrak{g} is of finite type, depending only on the derived Lie algebra \mathfrak{g}' and acting on \mathfrak{g} via the adjoint representation $\text{Ad} : G \rightarrow \text{Aut}(\mathfrak{g})$. It is generated by the one-parameter subgroups $U_\alpha = \exp(\mathfrak{g}_\alpha)$, $\alpha \in \Delta^{re}$, and $\text{Ad}(U_\alpha) = \exp(\text{ad}_{\mathfrak{g}_\alpha})$. In the definitions of J. Tits [22] G is the group of complex points of \mathfrak{G}_D where D is the datum associated to A and the \mathbb{Z} -dual Λ of $\bigoplus_{i \in I} \mathbb{Z}\alpha_i^\vee$.

The Cartan subalgebras of \mathfrak{g} are G -conjugate. If \mathfrak{g} is indecomposable and not of finite type, there are exactly two conjugate classes (under the adjoint action of G) of Borel subalgebras : $G.\mathfrak{b}^+$ and $G.\mathfrak{b}^-$. A Borel subalgebra \mathfrak{b} of \mathfrak{g} which is G -conjugate to \mathfrak{b}^+ (resp. \mathfrak{b}^-) is called positive (resp. negative). It follows that any parabolic subalgebra \mathfrak{p} of \mathfrak{g} is G -conjugate to a standard positive (or negative) parabolic subalgebra, in which case, we say that \mathfrak{p} is positive (or negative).

Standard Kac-Moody subalgebras and subgroups

Let J be a non-empty subset of I . Consider the generalized Cartan matrix $A_J = (a_{i,j})_{i,j \in J}$.

Definition 1.1. The subset J is called of finite type if the corresponding generalized Cartan matrix A_J is. We say also that J is connected, if the Dynkin subdiagram, with vertices indexed by J , is connected or, equivalently, the corresponding generalized Cartan submatrix A_J is indecomposable.

Proposition 1.2. Let $\Pi_J = \{\alpha_j, j \in J\}$ and $\Pi_{\check{J}} = \{\alpha_{\check{j}}, j \in J\}$. Let \mathfrak{h}'_J be the subspace of \mathfrak{h} generated by $\Pi_{\check{J}}$, and $\mathfrak{h}^J = \Pi_J^\perp = \{h \in \mathfrak{h}, \langle \alpha_j, h \rangle = 0, \forall j \in J\}$. Let \mathfrak{h}''_J be a supplementary subspace of $\mathfrak{h}'_J + \mathfrak{h}^J$ in \mathfrak{h} and let

$$\mathfrak{h}_J = \mathfrak{h}'_J \oplus \mathfrak{h}''_J,$$

then, we have :

- 1) $(\mathfrak{h}_J, \Pi_J, \Pi_{\check{J}})$ is a realization of the generalized Cartan matrix A_J . Hence $\mathfrak{h}''_J = \{0\}$, $\mathfrak{h}_J = \mathfrak{h}'_J$ when A_J is regular (e.g. when J is of finite type).
- 2) The subalgebra $\mathfrak{g}(J)$ of \mathfrak{g} , generated by \mathfrak{h}_J and the $e_j, f_j, j \in J$, is the Kac-Moody Lie algebra associated to the realization $(\mathfrak{h}_J, \Pi_J, \Pi_{\check{J}})$ of A_J .
- 3) The corresponding root system $\Delta(J) = \Delta(\mathfrak{g}(J), \mathfrak{h}_J)$ can be identified with $\Delta_J := \Delta \cap (\bigoplus_{j \in J} \mathbb{Z}\alpha_j)$.

N.B. The derived algebra $\mathfrak{g}'(J)$ of $\mathfrak{g}(J)$ is generated by the e_j, f_j for $j \in J$; it does not depend of the choice of \mathfrak{h}''_J .

Proof. We may assume \mathfrak{g} indecomposable.

1) Note that $\dim(\mathfrak{h}''_J) = \dim(\mathfrak{h}'_J \cap \mathfrak{h}^J) = \text{corank}(A_J)$. In particular, $\dim(\mathfrak{h}_J) - |J| = \text{corank}(A_J)$. If $\alpha \in \text{Vect}(\alpha_j, j \in J)$, then α is entirely determined by its restriction to \mathfrak{h}_J and hence Π_J defines, by restriction, a linearly independent

set in \mathfrak{h}_J^* . As Π_J is linearly independent, assertion 1) holds. Assertions 2) and 3) are straightforward. ■

In the same way, the subgroup G_J of G generated by $U_{\pm\alpha_j}$, $j \in J$, is equal to the Kac-Moody group associated to the generalized Cartan matrix A_J : it is clearly a quotient; the well known equality is proven explicitly in [20, 5.15.2], it may be deduced from [22, th. 1], see also [19, 8.4.2].

The invariant bilinear form (See [12]).

We recall that the generalized Cartan matrix A is supposed symmetrizable. There exists a non-degenerate $\text{ad}(\mathfrak{g})$ -invariant symmetric \mathbb{C} -bilinear form (\cdot, \cdot) on \mathfrak{g} , which is entirely determined by its restriction to \mathfrak{h} , such that

$$(\alpha_i, h) = \frac{(\alpha_i, \alpha_i)}{2} \langle \alpha_i, h \rangle, \quad i \in I, h \in \mathfrak{h},$$

and we may thus assume that

$$(\alpha_i, \alpha_i) \text{ is a positive rational for all } i. \tag{1.2}$$

The non-degenerate invariant bilinear form (\cdot, \cdot) induces an isomorphism $\nu : \mathfrak{h} \rightarrow \mathfrak{h}^*$ such that $\alpha_i = \frac{2\nu(\alpha_i)}{(\alpha_i, \alpha_i)}$ and $\alpha_i = \frac{2\nu^{-1}(\alpha_i)}{(\alpha_i, \alpha_i)}$ for all i .

There exists a totally isotropic subspace \mathfrak{h}'' of \mathfrak{h} (relative to the invariant bilinear form (\cdot, \cdot)) which is in duality with the center \mathfrak{c} of \mathfrak{g} . In particular, \mathfrak{h}'' defines a supplementary subspace of \mathfrak{h}' in \mathfrak{h} .

Note that any invariant symmetric bilinear form b on \mathfrak{g} satisfying $b(\alpha_i, \alpha_i) > 0$, $\forall i \in I$, is non-degenerate and $b(\alpha_i, h) = \frac{b(\alpha_i, \alpha_i)}{2} \langle \alpha_i, h \rangle$, $\forall i \in I, \forall h \in \mathfrak{h}$. It follows that, if \mathfrak{g} is indecomposable, the restriction of b to \mathfrak{g}' is proportional to that of (\cdot, \cdot) . In particular, if moreover A is non-singular, then the invariant bilinear form (\cdot, \cdot) satisfying the condition 1.2 is unique up to a positive rational factor.

The Tits cone (See [12, Chap. 3 and 5]).

Let $C := \{h \in \mathfrak{h}_{\mathbb{R}}; \langle \alpha_i, h \rangle \geq 0, \forall i \in I\}$ be the fundamental chamber (relative to the root basis Π) and let $X := \bigcup_{w \in W} w(C)$ be the Tits cone. We have the following

description of the Tits cone:

- (1) $X = \{h \in \mathfrak{h}_{\mathbb{R}}; \langle \alpha, h \rangle < 0 \text{ only for a finite number of } \alpha \in \Delta^+\}$.
- (2) $X = \mathfrak{h}_{\mathbb{R}}$ if and only if the generalized Cartan matrix A is of finite type.
- (3) If A is indecomposable of affine type, then $X = \{h \in \mathfrak{h}_{\mathbb{R}}; \langle \delta, h \rangle > 0\} \cup \mathbb{R}\nu^{-1}(\delta)$, where δ is the lowest imaginary positive root of Δ^+ .
- (4) If A is indecomposable of indefinite type, then the closure of the Tits cone, for the metric topology on $\mathfrak{h}_{\mathbb{R}}$, is $\bar{X} = \{h \in \mathfrak{h}_{\mathbb{R}}; \langle \alpha, h \rangle \geq 0, \forall \alpha \in \Delta_{im}^+\}$.
- (5) If $h \in X$, then h lies in the interior $\overset{\circ}{X}$ of X if and only if the fixer W_h of h , in the Weyl group W , is finite. Thus $\overset{\circ}{X}$ is the union of finite type facets of X .
- (6) If A is hyperbolic, then $\bar{X} \cup (-\bar{X}) = \{h \in \mathfrak{h}_{\mathbb{R}}; (h, h) \leq 0\}$ and the set of imaginary roots is $\Delta^{im} = \{\alpha \in Q \setminus \{0\}; (\alpha, \alpha) \leq 0\}$, where $Q = \mathbb{Z}\Pi$ is the root lattice.

Remark 1.3. Combining (3) and (4) one obtains that if A is not of finite type then $\bar{X} = \{h \in \mathfrak{h}_{\mathbb{R}}; \langle \alpha, h \rangle \geq 0, \forall \alpha \in \Delta_{im}^+\}$.

Graded Kac-Moody Lie algebras

Definition 1.4. Let Σ be a root system of Kac-Moody type. The Kac-Moody Lie algebra \mathfrak{g} is said to be finitely Σ -graded if :

(i) \mathfrak{g} contains, as a subalgebra, a Kac-Moody algebra \mathfrak{m} whose root system relative to a Cartan subalgebra \mathfrak{a} is equal to Σ .

(ii) $\mathfrak{g} = \sum_{\alpha \in \Sigma \cup \{0\}} V_{\alpha}$, with $V_{\alpha} = \{x \in \mathfrak{g}; [a, x] = \langle \alpha, a \rangle x, \forall a \in \mathfrak{a}\}$.

(iii) V_{α} is finite dimensional for all $\alpha \in \Sigma \cup \{0\}$.

We say that \mathfrak{m} (as in (i) above) is a grading subalgebra, and $(\mathfrak{g}, \Sigma, \mathfrak{m})$ a gradation with finite multiplicities (or, to be short, a finite gradation).

Note that from (ii) the Cartan subalgebra \mathfrak{a} of \mathfrak{m} is $\text{ad}(\mathfrak{g})$ -diagonalizable, and we may assume that \mathfrak{a} is contained in the standard Cartan subalgebra \mathfrak{h} of \mathfrak{g} .

Lemma 1.5. *Let \mathfrak{g} be a Kac-Moody algebra finitely Σ -graded, with grading subalgebra \mathfrak{m} . If \mathfrak{m} itself is finitely Σ' -graded (for some root system Σ' of Kac-Moody type), then \mathfrak{g} is finitely Σ' -graded.*

Proof. If \mathfrak{m}' is the grading subalgebra of \mathfrak{m} , we may suppose the Cartan subalgebras such that $\mathfrak{a}' \subset \mathfrak{a} \subset \mathfrak{h}$, with obvious notations. Conditions (i) and (ii) are clearly satisfied for \mathfrak{g} , \mathfrak{m}' and \mathfrak{a}' . Condition (iii) for \mathfrak{m} and Σ' tells that, for all $\alpha' \in \Sigma'$, the set $\{\alpha \in \Sigma \mid \alpha|_{\mathfrak{a}'} = \alpha'\}$ is finite. But $V_{\alpha'} = \bigoplus_{\alpha|_{\mathfrak{a}'} = \alpha'} V_{\alpha}$, so each $V_{\alpha'}$ is finite dimensional if this is true for each V_{α} . ■

Examples 1.6.

1) Let $\Delta = \Delta(\mathfrak{g}, \mathfrak{h})$ the root system of \mathfrak{g} relative to \mathfrak{h} , then \mathfrak{g} is finitely Δ -graded: this is the trivial gradation of \mathfrak{g} by its own root system.

2) Let $\mathfrak{g}_{\mathbb{R}}$ be an almost split real form of \mathfrak{g} (see [2]) and let $\mathfrak{t}_{\mathbb{R}}$ be a maximal split toral subalgebra of $\mathfrak{g}_{\mathbb{R}}$. Suppose that the restricted root system $\Delta' = \Delta(\mathfrak{g}_{\mathbb{R}}, \mathfrak{t}_{\mathbb{R}})$ is reduced of Kac-Moody type. In [4, §9], N. Bardy constructed a split real Kac-Moody subalgebra $\mathfrak{l}_{\mathbb{R}}$ of $\mathfrak{g}_{\mathbb{R}}$ such that $\Delta' = \Delta(\mathfrak{l}_{\mathbb{R}}, \mathfrak{t}_{\mathbb{R}})$, then \mathfrak{g} is obviously finitely Δ' -graded.

We get thus many examples coming from known tables for almost split real forms: see [2] in the affine case and [6] in the hyperbolic case.

3) When $\mathfrak{g}_{\mathbb{R}}$ is an almost compact real form of \mathfrak{g} , the same constructions should lead to gradations by finite root systems, as in [5] e.g.

2. Gradations associated to C -admissible pairs

In this section, we suppose the Kac-Moody Lie algebra \mathfrak{g} indecomposable and symmetrizable, see however Remark 2.15. We shall build a finite gradation of \mathfrak{g} associated to some good subset of I .

We recall some definitions introduced by H. Rubenthaler ([21]) and J. Nervi ([16], [17]). Let J be a subset of I of finite type. For $k \in I \setminus J$, we denote by I_k the connected component, containing k , of the Dynkin subdiagram corresponding to $J \cup \{k\}$, and let $J_k := I_k \setminus \{k\}$.

We are interested in the case where I_k is of finite type for all $k \in I \setminus J$: that is always true if \mathfrak{g} is of affine type and $|I \setminus J| \geq 2$ or if \mathfrak{g} is of hyperbolic type and $|I \setminus J| \geq 3$.

For $k \in I \setminus J$, let $\mathfrak{g}(I_k)$ be the simple subalgebra generated by $\mathfrak{g}_{\pm\alpha_i}$, $i \in I_k$, then $\mathfrak{h}_{I_k} = \mathfrak{h} \cap \mathfrak{g}(I_k) = \sum_{i \in I_k} \mathbb{C}\alpha_i$ is a Cartan subalgebra of $\mathfrak{g}(I_k)$. Let H_k be the unique element of \mathfrak{h}_{I_k} such that $\langle \alpha_i, H_k \rangle = 2\delta_{i,k}$, $\forall i \in I_k$.

Definition 2.1. We suppose the Dynkin diagram indexed by I connected and consider a subset J of finite type. We preserve the notations introduced above.

- 1) Let $k \in I \setminus J$.
 - (i) The pair (I_k, J_k) is called admissible if I_k is of finite type and there exist $E_k, F_k \in \mathfrak{g}(I_k)$ such that (E_k, H_k, F_k) is an \mathfrak{sl}_2 -triple.
 - (ii) The pair (I_k, J_k) is called C -admissible if it is admissible and the simple Lie algebra $\mathfrak{g}(I_k)$ is A_1 -graded by the root system, of type A_1 , associated to the \mathfrak{sl}_2 -triple (E_k, H_k, F_k) .
- 2) The pair (I, J) is called C -admissible if the pairs (I_k, J_k) are C -admissible for all $k \in I \setminus J$. It is said irreducible if, moreover, $|I \setminus J| = 1$.

Schematically, any C -admissible pair (I, J) is represented by the Dynkin diagram, corresponding to A , on which the vertices indexed by J are denoted by white circles \circ and those of $I \setminus J$ are denoted by black circles \bullet .

Remark 2.2. 1) The admissibility of each (I_k, J_k) is essential to build (in 2.6, 2.11) the grading subalgebra \mathfrak{g}^J and its grading root system Δ^J .

2) As $\mathfrak{g}(J)$ will be in the eigenspace V_0 of weight 0 for the grading by Δ^J , it is necessary to assume J of finite type to get a finite gradation.

3) I_k is of finite type if, and only if, $\mathfrak{g}(I_k)$ is finite dimensional, and this is equivalent to the alternative assumption in (ii) that the A_1 -gradation has finite multiplicities. It is clearly necessary to get, in Theorem 2.14, a finite gradation of \mathfrak{g} by the root system Δ^J . Moreover, even in a more general situation, the condition I_k of finite type will naturally appear (3.14).

4) Note that the definition presented here, for C -admissible pairs, is equivalent to that introduced by Rubenthaler and Nervi (see [21], [16]) in terms of prehomogeneous spaces of parabolic type : if (I_k, J_k) is C -admissible, define for $p \in \mathbb{Z}$, the subspace $d_{k,p} := \{X \in \mathfrak{g}(I_k); [H_k, X] = 2pX\}$; then $(d_{k,0}, d_{k,1})$ is an irreducible regular and commutative prehomogeneous space of parabolic type, and $d_{k,p} = \{0\}$ for $|p| \geq 2$. Then (I_k, J_k) is an irreducible C -admissible pair. According to Rubenthaler and Nervi ([21, Table 1] or [16, Table 2]) the irreducible C -admissible pair (I_k, J_k) should be among the list in Table 1 below.

5) Along our study of general finite gradations in section 3, we shall meet a situation of "generalized C -admissible pair" (I, J) (3.16) where $J \subset I$ is of finite type and I_k (for $k \in I' = I \setminus J$) is defined as above but perhaps not of

finite type. When k is in some subset I'_{re} of I' , (I_k, J_k) is C -admissible and the $k \in I'_{im} = I' \setminus I'_{re}$ do not contribute to the root system Σ grading \mathfrak{g} . But we do not know the good assumptions on these (I_k, J_k) for $k \in I'_{im}$ to get, conversely, a finite gradation of \mathfrak{g} by some root system. So we give no precise definition; it is expected in the work in preparation [7].

Table 1

List of irreducible C -admissible pairs

$A_{2n-1}, n \geq 1$	
$B_n, n \geq 3$	
$C_n, n \geq 2$	
$D_{n,1}, n \geq 4$	
$D_{2n,2}, n \geq 2$	
E_7	

Definition 2.3. Let J be a subset of I and let $i, k \in I \setminus J$. We say that i and k are J -connected relative to A if there exist $j_0, j_1, \dots, j_{p+1} \in I$ such that $j_0 = i, j_{p+1} = k, j_s \in J, \forall s = 1, 2, \dots, p$, and $a_{j_s, j_{s+1}} \neq 0, \forall s = 0, 1, \dots, p$.

Remark 2.4. Note that the relation “to be J -connected” is symmetric on i and k . As the generalized Cartan matrix A is assumed to be indecomposable, for any vertices $i, k \in I \setminus J$ there exist $i_0, i_1, \dots, i_{p+1} \in I \setminus J$ such that $i_0 = i, i_{p+1} = k$ and i_s and i_{s+1} are J -connected for all $s = 0, 1, \dots, p$.

Let us assume from now on that (I, J) is a C -admissible pair and let $I' := I \setminus J$. For $k \in I'$, let (E_k, H_k, F_k) be an \mathfrak{sl}_2 -triple associated to the irreducible C -admissible pair (I_k, J_k) .

Lemma 2.5. *Let $k \neq l \in I'$, then :*

- 1) $\langle \alpha_l, H_k \rangle \in \mathbb{Z}^-$.
- 2) *the following assertions are equivalent :*
 - i) k, l are J -connected
 - ii) $\langle \alpha_l, H_k \rangle$ is a negative integer
 - iii) $\langle \alpha_k, H_l \rangle$ is a negative integer

Proof. 1) One can write $H_k = \sum_{i \in I_k} n_{i,k} \alpha_{\check{i}}$, where $n_{i,k}$ are positive integers (see [21] or [17, 1.4.1.2]). As $l \notin I_k$, we have that $\langle \alpha_l, H_k \rangle = \sum_{i \in I_k} n_{i,k} \langle \alpha_l, \alpha_{\check{i}} \rangle \in \mathbb{Z}^-$.

2) In view of Remark 2.4, it suffices to prove the equivalence between i) and ii). Since I_k is the connected component of $J \cup \{k\}$ containing k , the assertion i) is equivalent to say that the vertex l is connected to I_k , so there exists $i_k \in I_k$ such that $\langle \alpha_l, \alpha_{\check{i}_k} \rangle < 0$ and hence $\langle \alpha_l, H_k \rangle < 0$. ■

Proposition 2.6. *Let $\mathfrak{h}^J = \Pi_J^\perp = \{h \in \mathfrak{h}, \langle \alpha_j, h \rangle = 0, \forall j \in J\}$. For $k \in I'$, denote by $\alpha'_k = \alpha_k / \mathfrak{h}^J$ the restriction of α_k to the subspace \mathfrak{h}^J of \mathfrak{h} , and $\Pi^J = \{\alpha'_k; k \in I'\}$, $\Pi^{J\vee} = \{H_k; k \in I'\}$. For $k, l \in I'$, put $a'_{k,l} = \langle \alpha_l, H_k \rangle$ and $A^J = (a'_{k,l})_{k,l \in I'}$. Then A^J is an indecomposable and symmetrizable generalized Cartan matrix, $(\mathfrak{h}^J, \Pi^J, \Pi^{J\vee})$ is a realization of A^J and $\text{corank}(A^J) = \text{corank}(A)$.*

Proof. The fact that $a'_{k,k} = 2$ follows from the definition of H_k for $k \in I'$. If $k \neq l \in I'$, then by lemma 2.5, $a'_{k,l} \in \mathbb{Z}^-$ and $a'_{k,l} \neq 0$ if and only if $a'_{l,k} \neq 0$. Hence A^J is a generalized Cartan matrix. As the matrix A is indecomposable, A_J is also indecomposable (see Remark 2.4). Clearly $\Pi^J = \{\alpha'_k; k \in I'\}$ is a linearly independent subset of the dual space \mathfrak{h}^{J*} of \mathfrak{h}^J , $\Pi^{J\vee} = \{H_k; k \in I'\}$ is a linearly independent subset of \mathfrak{h}^J and by construction $\langle \alpha_l, H_k \rangle = a'_{k,l}, \forall k, l \in I'$. We have to prove that $\dim(\mathfrak{h}^J) - |I'| = \text{corank}(A^J)$. As J is of finite type, the restriction of the invariant bilinear form (\cdot, \cdot) to \mathfrak{h}_J is non-degenerate and \mathfrak{h}_J is contained in $\mathfrak{h}' = \bigoplus_{i \in I} \mathbb{C}\alpha_{\check{i}}$. Therefore

$$\mathfrak{h} = \mathfrak{h}^J \oplus^\perp \mathfrak{h}_J$$

and

$$\mathfrak{h}' = (\mathfrak{h}' \cap \mathfrak{h}^J) \oplus \mathfrak{h}_J.$$

It follows that $\dim(\mathfrak{h}' \cap \mathfrak{h}^J) = |I'| = \dim(\bigoplus_{k \in I'} \mathbb{C}H_k)$. As the subspace $\bigoplus_{k \in I'} \mathbb{C}H_k$ is contained in $\mathfrak{h}' \cap \mathfrak{h}^J$, we deduce that $\mathfrak{h}' \cap \mathfrak{h}^J = \bigoplus_{k \in I'} \mathbb{C}H_k$. Note that any supplementary subspace $\mathfrak{h}^{J''}$ of $\mathfrak{h}' \cap \mathfrak{h}^J$ in \mathfrak{h}^J is also a supplementary of \mathfrak{h}' in \mathfrak{h} ; hence, we have that $\text{corank}(A) = \dim(\mathfrak{h}^{J''}) = \dim(\mathfrak{h}^J) - |I'|$. Let $\mathfrak{c} := \bigcap_{i \in I} \ker(\alpha_i)$ be the center of \mathfrak{g} and let $\mathfrak{c}^J = \bigcap_{k \in I'} \ker(\alpha'_k)$. Recall that $\text{corank}(A) = \dim(\mathfrak{c})$ and $\text{corank}(A^J) = \dim(\mathfrak{c}^J)$. It's clear that $\mathfrak{c}^J = \mathfrak{c}$; hence $\text{corank}(A^J) = \dim(\mathfrak{c}^J) = \text{corank}(A) = \dim(\mathfrak{h}^J) - |I'|$.

It remains to prove that A^J is symmetrizable. For $k \in I'$, let R_k^J be the fundamental reflection of \mathfrak{h}^J such that $R_k^J(h) = h - \langle \alpha'_k, h \rangle H_k, \forall h \in \mathfrak{h}^J$. Let W^J be the Weyl group of A^J generated by $R_k^J, k \in I'$. Let $(\cdot, \cdot)^J$ be the restriction

to \mathfrak{h}^J of the invariant bilinear form (\cdot, \cdot) on \mathfrak{h} . Then $(\cdot, \cdot)^J$ is a non-degenerate symmetric bilinear form on \mathfrak{h}^J which is W^J -invariant (see the lemma hereafter). From the relation $(R_k^J(H_k), R_k^J(H_l))^J = (H_k, H_l)^J$ one can deduce that

$$(H_k, H_l)^J = \frac{(H_k, H_k)^J}{2} a'_{l,k}, \forall k, l \in I'.$$

Since $(H_k, H_k)^J > 0, \forall k \in I'$, the generalized Cartan matrix ${}^tA^J$ (and so A^J) is symmetrizable. ■

Lemma 2.7. For $k \in I' := I \setminus J$, let w_k^J be the longest element of the Weyl group $W(I_k)$ generated by the fundamental reflections $r_i, i \in I_k$. Then w_k^J stabilizes \mathfrak{h}^J and induces the fundamental reflection R_k^J of \mathfrak{h}^J associated to H_k .


Proof. If one looks at the list above of the irreducible C -admissible pairs, one can see that $w_k^J(\alpha_k) = -\alpha_k$ and that $-w_k^J$ permutes the $\alpha_j, j \in J_k$. Clearly $w_k^J(\alpha_j) = \alpha_j, \forall j \in J \setminus J_k$. Hence w_k^J stabilizes \mathfrak{h}_J and its orthogonal subspace $\mathfrak{h}_J^\perp = \mathfrak{h}^J$. Note that $-w_k^J(H_k) \in \mathfrak{h}_{I_k}$ and it satisfies the same equations defining H_k . Hence $-w_k^J(H_k) = H_k = -R_k^J(H_k)$. Recall that $\ker(\alpha'_k) = \ker(\alpha_k) \cap (\bigcap_{j \in J} \ker(\alpha_j))$; thus it is fixed by R_k^J and W_k^J . Since $\mathfrak{h}^J = \ker(\alpha'_k) \oplus \mathbb{C}H_k$, the reflection R_k^J coincides with W_k^J on \mathfrak{h}^J . ■

Remark 2.8. Actually we can now rediscover the list of irreducible C -admissible pairs given in Table 1. The black vertex k should be invariant under $-w_k^J$ and the corresponding coefficient of the highest root of I_k should be 1 (an easy consequence of the definition 2.1 1) (ii)).


Example 2.9. Consider the hyperbolic generalized Cartan matrix A of type $HE_8^{(1)} = E_{10}$ indexed by $I = \{-1, 0, 1, \dots, 8\}$.

The following two choices for J define C -admissible pairs :


1) $J = \{2, 3, 4, 5\}$.



The corresponding generalized Cartan matrix A^J is hyperbolic of type $HF_4^{(1)}$:



2) $J = \{1, 2, 3, 4, 5, 6\}$.



The corresponding generalized Cartan matrix A^J is hyperbolic of type $HG_2^{(1)}$:



Note that the first example corresponds to an almost split real form of the Kac-Moody Lie algebra $\mathfrak{g}(A)$ and A^J is the generalized Cartan matrix associated to the corresponding (reduced) restricted root system (see [6]) whereas the second example does not correspond to an almost split real form of $\mathfrak{g}(A)$.

Lemma 2.10. For $k \in I'$, set $\mathfrak{s}(k) = \mathbb{C}E_k \oplus \mathbb{C}H_k \oplus \mathbb{C}F_k$. Then, the Lie algebra \mathfrak{g} is an integrable $\mathfrak{s}(k)$ -module via the adjoint representation of $\mathfrak{s}(k)$ on \mathfrak{g} .

Proof. Note that $\mathfrak{s}(k)$ is isomorphic to $\mathfrak{sl}_2(\mathbb{C})$ with standard basis (E_k, H_k, F_k) . It is clear that $\text{ad}(H_k)$ is diagonalizable on \mathfrak{g} and $E_k = \sum_{\alpha} e_{\alpha} \in d_{k,1}$, where α runs over the set $\Delta_{k,1} = \{\alpha \in \Delta(I_k); \langle \alpha, H_k \rangle = 2\}$, $e_{\alpha} \in \mathfrak{g}_{\alpha}$ for $\alpha \in \Delta(I_k)$, and $d_{k,1} := \{X \in \mathfrak{g}(I_k); [H_k, X] = 2X\}$. Since $\Delta_{k,1} \subset \Delta^{re}$, $\text{ad}(e_{\alpha})$ is locally nilpotent for $\alpha \in \Delta_{k,1}$. As $d_{k,1}$ is commutative (see Remark 2.2) we deduce that $\text{ad}(E_k)$ is locally nilpotent on \mathfrak{g} . The same argument shows that $\text{ad}(F_k)$ is also locally nilpotent. Hence, the Kac-Moody algebra \mathfrak{g} is an integrable $\mathfrak{s}(k)$ -module. \blacksquare

Proposition 2.11. *Let \mathfrak{g}^J be the subalgebra of \mathfrak{g} generated by \mathfrak{h}^J and $E_k, F_k, k \in I'$. Then \mathfrak{g}^J is the Kac-Moody Lie algebra associated to the realization $(\mathfrak{h}^J, \Pi^J, \Pi^{J\vee})$ of the generalized Cartan matrix A^J .*

Proof. It is not difficult to check that the following relations hold in the Lie subalgebra \mathfrak{g}^J :

$$\begin{aligned} [\mathfrak{h}^J, \mathfrak{h}^J] &= 0, & [E_k, F_l] &= \delta_{k,l}H_k & (k, l \in I'); \\ [h, E_k] &= \langle \alpha'_k, h \rangle E_k, & [h, F_k] &= -\langle \alpha'_k, h \rangle F_k & (h \in \mathfrak{h}^J, k \in I'). \end{aligned}$$

We have to prove the Serre's relations :

$$(\text{ad}E_k)^{1-a'_{k,l}}(E_l) = 0, \quad (\text{ad}F_k)^{1-a'_{k,l}}(F_l) = 0 \quad (k \neq l \in I').$$

For $k \in I'$, let $\mathfrak{s}(k) = \mathbb{C}F_k \oplus \mathbb{C}H_k \oplus \mathbb{C}E_k$ be the Lie subalgebra of \mathfrak{g} isomorphic to $\mathfrak{sl}_2(\mathbb{C})$. Let $l \neq k \in I'$; note that $[H_k, F_l] = -a'_{k,l}F_l$ and $[E_k, F_l] = 0$, which means that F_l is a primitive weight vector for $\mathfrak{s}(k)$. As \mathfrak{g} is an integrable $\mathfrak{s}(k)$ -module (see Lemma 2.10) the primitive weight vector F_l is contained in a finite dimensional $\mathfrak{s}(k)$ -submodule (see [12, 3.6]). The relation $(\text{ad}F_k)^{1-a'_{k,l}}(F_l) = 0$ follows from the representation theory of $\mathfrak{sl}_2(\mathbb{C})$ (see [12, 3.2]). By similar arguments we prove that $(\text{ad}E_k)^{1-a'_{k,l}}(E_l) = 0$.

Now \mathfrak{g}^J is a quotient of the Kac-Moody algebra associated to A^J and $(\mathfrak{h}^J, \Pi^J, \Pi^{J\vee})$. By [12, 1.7] it is equal to it. \blacksquare

Definition 2.12. The Kac-Moody Lie algebra \mathfrak{g}^J is called the C -admissible algebra associated to the C -admissible pair (I, J) .

Proposition 2.13. *The Kac-Moody algebra \mathfrak{g} is an integrable \mathfrak{g}^J -module with finite multiplicities.*

Proof. The \mathfrak{g}^J -module \mathfrak{g} is clearly $\text{ad}(\mathfrak{h}^J)$ -diagonalizable and $\text{ad}(E_k), \text{ad}(F_k)$ are locally nilpotent on \mathfrak{g} for $k \in I'$ (see Lemma 2.10). Hence, \mathfrak{g} is an integrable \mathfrak{g}^J -module. For $\alpha \in \Delta$, let $\alpha' = \alpha|_{\mathfrak{h}^J}$ be the restriction of α to \mathfrak{h}^J . Set $\Delta' = \{\alpha'; \alpha \in \Delta\} \setminus \{0\}$. Then the set of weights, for the \mathfrak{g}^J -module \mathfrak{g} , is exactly $\Delta' \cup \{0\}$. Note that for $\alpha \in \Delta$, $\alpha' = 0$ if and only if $\alpha \in \Delta(J)$. In particular, the weight space $V_0 = \mathfrak{h} \oplus (\bigoplus_{\alpha \in \Delta(J)} \mathfrak{g}_{\alpha})$ corresponding to the null weight is finite dimensional. Let $\alpha = \sum_{i \in I} n_i \alpha_i \in \Delta$ such that $\alpha' \neq 0$. We will see that the corresponding weight space $V_{\alpha'}$ is finite dimensional. Note that $V_{\alpha'} = \bigoplus_{\beta' = \alpha'} \mathfrak{g}_{\beta}$. Let $\beta = \sum_{i \in I} m_i \alpha_i \in \Delta$ such that $\beta' = \alpha' = \sum_{k \in I'} n_k \alpha'_k$, then $m_k = n_k, \forall k \in I'$, since $\Pi^J = \{\alpha'_k, k \in I'\}$ is free in $(\mathfrak{h}^J)^*$. In particular, β and α are of the same

sign, and we may assume $\alpha \in \Delta^+$. Let $ht_J(\beta) = \sum_{j \in J} m_j$ be the height of β relative to J , and let W_J be the finite subgroup of W generated by $r_j, j \in J$. Since W_J fixes pointwise \mathfrak{h}^J , we deduce that $\gamma' = \beta', \forall \gamma \in W_J\beta$, and so we may assume that $ht_J(\beta)$ is minimal among the roots in $W_J\beta$. From the inequality $ht_J(\beta) \leq ht_J(r_j(\beta)), \forall j \in J$, we get $\langle \beta, \alpha_{\check{j}} \rangle \leq 0, \forall j \in J$. Let $\rho_{\check{J}}$ be the half sum of positive coroots of $\Delta(J)$. It is known that $\langle \alpha_j, \rho_{\check{J}} \rangle = 1, \forall j \in J$. Note that $\langle \beta, \rho_{\check{J}} \rangle = \sum_{j \in J} m_j + \sum_{k \in I'} n_k \langle \alpha_k, \rho_{\check{J}} \rangle = ht_J(\beta) + \sum_{k \in I'} n_k \langle \alpha_k, \rho_{\check{J}} \rangle$. Hence, the condition $(\langle \beta, \rho_{\check{J}} \rangle \leq 0)$ implies $(ht_J(\beta) \leq \sum_{k \in I'} -n_k \langle \alpha_k, \rho_{\check{J}} \rangle)$. Thus there is just a finite number of possibilities for β . It follows that α' is of finite multiplicity. ■

Theorem 2.14. *Let Δ^J be the root system of the pair $(\mathfrak{g}^J, \mathfrak{h}^J)$, then the Kac-Moody Lie algebra \mathfrak{g} is finitely Δ^J -graded, with grading subalgebra \mathfrak{g}^J .*

Proof. Let $\Delta' = \{\alpha', \alpha \in \Delta\} \setminus \{0\}$ be the set of non-null weights of the \mathfrak{g}^J -module \mathfrak{g} relative to \mathfrak{h}^J . Let $\Delta'_+ = \{\alpha' \in \Delta', \alpha \in \Delta^+\}$ and Δ^J_+ the set of positive roots of Δ^J relative to the root basis Π^J . We have to prove that $\Delta' = \Delta^J$ or equivalently $\Delta'_+ = \Delta^J_+$. Let $Q^J = \mathbb{Z}\Pi^J$ be the root lattice of Δ^J and $Q^J_+ = \mathbb{Z}^+\Pi^J$. It is known that the positive root system Δ^J_+ is uniquely defined by the following properties (see [12, Ex. 5.4]) :

- (i) $\Pi^J \subset \Delta^J_+ \subset Q^J_+, 2\alpha'_i \notin \Delta^J_+, \forall i \in I'$;
- (ii) if $\alpha' \in \Delta^J_+, \alpha' \neq \alpha'_i$, then the set $\{\alpha' + k\alpha'_i; k \in \mathbb{Z}\} \cap \Delta^J_+$ is a string $\{\alpha' - p\alpha'_i, \dots, \alpha' + q\alpha'_i\}$, where $p, q \in \mathbb{Z}^+$ and $p - q = \langle \alpha', H_i \rangle$;
- (iii) if $\alpha' \in \Delta^J_+$, then $\text{supp}(\alpha')$ is connected.

We will see that Δ'_+ satisfies these three properties and hence $\Delta'_+ = \Delta^J_+$. Clearly $\Pi^J \subset \Delta'_+ \subset Q^J_+$. For $\alpha \in \Delta$ and $k \in I'$, the condition $\alpha' \in \mathbb{N}\alpha_k$ implies $\alpha \in \Delta(I_k)^+$. As (I_k, J_k) is C -admissible for $k \in I'$, the highest root of $\Delta(I_k)^+$ has coefficient 1 on the root α_k (cf. Remark 2.8). It follows that $2\alpha'_k \notin \Delta'_+$ and (i) is satisfied. By Proposition 2.13, \mathfrak{g} is an integrable \mathfrak{g}^J -module with finite multiplicities. Hence, the propriety (ii) follows from [12, 3.6]. Let $\alpha \in \Delta_+$, then $\text{supp}(\alpha)$ is connected and $\text{supp}(\alpha') \subset \text{supp}(\alpha)$. Let $k, l \in \text{supp}(\alpha')$; if k, l are J -connected in $\text{supp}(\alpha)$ relative to the generalized Cartan matrix A (cf. 2.3), then by lemma 2.5, k, l are linked in I' relative to the generalized Cartan matrix A^J . Hence, the connectedness of $\text{supp}(\alpha')$, relative to A^J , follows from that of $\text{supp}(\alpha)$ relative to A (see Remark 2.4) and (iii) is satisfied. ■

Remark 2.15. Note that the definition of C -admissible pair can be extended to decomposable Kac-Moody Lie algebras : thus if I^1, I^2, \dots, I^m are the connected components of I and $J^k = J \cap I^k, k = 1, 2, \dots, m$, then (I, J) is C -admissible if and only if (I^k, J^k) is for all $k = 1, 2, \dots, m$. In particular, the corresponding C -admissible algebra is $\mathfrak{g}^J = \bigoplus_{k=1}^m \mathfrak{g}(I^k)^{J^k}$, where $\mathfrak{g}(I^k)^{J^k}$ is the C -admissible subalgebra of $\mathfrak{g}(I^k)$ corresponding to the C -admissible pair $(I^k, J^k), k = 1, 2, \dots, m$.

3. Real gradations

From now on we suppose that the Kac-Moody Lie algebra \mathfrak{g} is symmetrizable and, starting from 3.5, indecomposable.

Let \mathfrak{m} be a Kac-Moody subalgebra of \mathfrak{g} and let \mathfrak{a} be a Cartan subalgebra of \mathfrak{m} . Put $\Sigma = \Delta(\mathfrak{m}, \mathfrak{a})$ the corresponding root system. We assume that $\mathfrak{a} \subset \mathfrak{h}$ and that \mathfrak{g} is finitely Σ -graded with \mathfrak{m} as grading subalgebra. Thus $\mathfrak{g} = \sum_{\gamma \in \Sigma \cup \{0\}} V_\gamma$, with $V_\gamma = \{x \in \mathfrak{g}; [a, x] = \langle \gamma, a \rangle x, \forall a \in \mathfrak{a}\}$ is finite dimensional for all $\gamma \in \Sigma \cup \{0\}$. For $\alpha \in \Delta$, denote by $\rho_\alpha(\alpha)$ the restriction of α to \mathfrak{a} . As \mathfrak{g} is Σ -graded, one has $\rho_\alpha(\Delta \cup \{0\}) = \Sigma \cup \{0\}$.

Lemma 3.1.

- 1) Let \mathfrak{c} be the center of \mathfrak{g} and denote by \mathfrak{c}_a the center of \mathfrak{m} . Then $\mathfrak{c}_a = \mathfrak{c} \cap \mathfrak{a}$. In particular, if \mathfrak{g} is perfect, then the grading subalgebra \mathfrak{m} is also perfect.
- 2) Suppose that $\Delta^{im} \neq \emptyset$, then $\rho_\alpha(\Delta^{im}) \subset \Sigma^{im}$.

Proof.

- 1) It is clear that $\mathfrak{c} \cap \mathfrak{a} \subset \mathfrak{c}_a$. Since \mathfrak{g} is Σ -graded, we deduce that \mathfrak{c}_a is contained in the center \mathfrak{c} of \mathfrak{g} , hence $\mathfrak{c}_a \subset \mathfrak{c} \cap \mathfrak{a}$. If \mathfrak{g} is perfect, then $\mathfrak{g} = \mathfrak{g}'$, $\mathfrak{h} = \mathfrak{h}'$, $\mathfrak{c} = \{0\}$; so $\mathfrak{c}_a = \{0\}$, $\mathfrak{a} = \mathfrak{a}'$ and $\mathfrak{m} = \mathfrak{m}'$.
- 2) If $\alpha \in \Delta^{im}$, then $\mathbb{N}\alpha \subset \Delta$. Since V_0 is finite dimensional, $\rho_\alpha(\alpha) \neq 0$ and $\mathbb{N}\rho_\alpha(\alpha) \subset \Sigma$, hence $\rho_\alpha(\alpha) \in \Sigma^{im}$. ■

Definition 3.2. ([3, 5.2.6]) Suppose that $\Delta^{im} \neq \emptyset$. Let $\alpha, \beta \in \Delta^{im}$.

- (i) The imaginary roots α and β are said to be linked if $\mathbb{N}\alpha + \mathbb{N}\beta \subset \Delta$ or $\beta \in \mathbb{Q}^+\alpha$.
- (ii) The imaginary roots α and β are said to be linkable if there exists a finite family of imaginary roots $(\beta_i)_{0 \leq i \leq n+1}$ such that $\beta_0 = \alpha$, $\beta_{n+1} = \beta$ and β_i and β_{i+1} are linked for all $i = 0, 1, \dots, n$.

Proposition 3.3. ([3, 5.2.7]) Suppose that $\Delta^{im} \neq \emptyset$. Let $\Delta = \bigcup_{j=1}^m \Delta_j$ be the decomposition of Δ in indecomposable root systems. Suppose that $\Delta_1, \Delta_2, \dots, \Delta_r$ ($r \leq m$) are the indecomposable root subsystems of Δ which are not of finite type. Then to be linkable is an equivalence relation on Δ^{im} and the equivalence classes are the $2r$ sets $\Delta_\pm^{im} \cap \Delta_j$, $j = 1, 2, \dots, r$.

Lemma 3.4. Suppose that $\Delta^{im} \neq \emptyset$, then there exist root bases in Σ and Δ such that $\rho_\alpha(\Delta_+^{im}) \subset \Sigma_+^{im}$.

Proof. Fix a root basis Π_a for the grading root system Σ . Let $\Delta = \bigcup_{j=1}^m \Delta_j$ be, as above, the decomposition of Δ in indecomposable root systems. Denote by $\Pi_j := \Pi \cap \Delta_j$ the root basis of Δ_j , $j = 1, 2, \dots, m$. If α, β are two imaginary linkable roots of Δ_j^{im} , then $\rho_\alpha(\alpha)$ and $\rho_\alpha(\beta)$ are also linkable in Σ^{im} . By Proposition 3.3, $\rho_\alpha(\alpha)$ and $\rho_\alpha(\beta)$ are of the same sign. Since α and β are of the same sign in Δ_j^{im} relative to the root basis Π_j , one can, if necessary, change the sign of Π_j so that $\rho_\alpha(\alpha)$ and $\rho_\alpha(\beta)$ are positive imaginary roots of Σ^+ relative to the fixed root basis

Π_a . Hence we get a root basis of $\Delta = \bigcup_{j=1}^m \Delta_j$ satisfying $\rho_a(\Delta_+^{im}) \subset \Sigma_+^{im}$. ■

In the following, we will show that the indecomposable Kac-Moody Lie algebra \mathfrak{g} and the grading subalgebra \mathfrak{m} are of the same type.

Lemma 3.5. *The Kac-Moody Lie algebra \mathfrak{g} is of indefinite type if and only if Δ^{im} generates the dual space $(\mathfrak{h}/\mathfrak{c})^*$ of $\mathfrak{h}/\mathfrak{c}$.*

Proof. Note that the root basis $\Pi = \{\alpha_i, i \in I\}$ induces a basis for the quotient vector space $(\mathfrak{h}/\mathfrak{c})^*$. It follows that the condition $(\Delta^{im} \neq \emptyset)$ implies $(\dim(\mathfrak{h}/\mathfrak{c})^* \geq 2)$. Suppose now that \mathfrak{g} is of indefinite type. Let $\alpha \in \Delta_+^{sim}$ be a positive strictly imaginary root satisfying $\langle \alpha, \alpha_i \rangle < 0, \forall i \in I$; then, $r_i(\alpha) = \alpha - \langle \alpha, \alpha_i \rangle \alpha_i \in \Delta_+^{im}$ for all $i \in I$. In particular, the vector subspace $\langle \Delta^{im} \rangle$ spanned by Δ^{im} contains Π and hence is equal to $(\mathfrak{h}/\mathfrak{c})^*$. Conversely, if Δ^{im} generates $(\mathfrak{h}/\mathfrak{c})^*$, then Δ^{im} is non-empty and contains at least two linearly independent imaginary roots; hence Δ can not be of finite or affine type. ■

Proposition 3.6.

- 1) If Δ^{im} is not empty, then \mathfrak{m} is indecomposable.
- 2) The Kac-Moody algebra \mathfrak{g} and the grading subalgebra \mathfrak{m} are of the same type.
- 3) Suppose \mathfrak{g} Lorentzian, then \mathfrak{m} is also Lorentzian.

N.B. We will see below that \mathfrak{m} is always indecomposable (3.11) and symmetrizable (3.17).

Proof.

1) We saw in Lemma 3.4 that $\rho_a(\Delta_+^{im})$ is in a unique linkable equivalence class of Σ_+^{im} . So, if $\Sigma = \Sigma_1 \cup \Sigma_2$ is decomposable, we may assume $\rho_a(\Delta_+^{im}) \subset \Sigma_1^{im}$. But there is $\delta \in \Delta_+^{im}$ such that $\alpha + n\delta \in \Delta_+$ for all $\alpha \in \Delta_+$ and $n \in \mathbb{N}$ [12, 4.3, 5.6 and 6.3]. So $\rho_a(\alpha) + n\rho_a(\delta) \in \Sigma$ for $n \gg 0$ and $\rho_a(\alpha) \in \Sigma_1 \cup \{0\}$. As $\rho_a(\Delta \cup \{0\}) = \Sigma \cup \{0\}$, we have $\Sigma_2 = \emptyset$.

2) If \mathfrak{g} is of finite type, then Δ is finite and hence $\Sigma = \rho_a(\Delta) \setminus \{0\}$ is finite. If \mathfrak{g} is affine, let δ be the lowest positive imaginary root. One can choose a root basis $\Pi_a = \{\gamma_i, i \in \bar{I}\}$ of Σ so that $\bar{\delta} := \rho_a(\delta)$ is a positive imaginary root. Note that $\mathfrak{a}' := \mathfrak{a} \cap \mathfrak{m}' \subset \mathfrak{h}'$; in particular $\bar{\delta}(\mathfrak{a}') = \{0\}$ and $\langle \bar{\delta}, \gamma_i \rangle = 0, \forall i \in \bar{I}$. It follows that \mathfrak{m} is affine (see [12, 4.3]).

Suppose now that \mathfrak{g} is of indefinite type. Thanks to Lemma 3.5, it suffices to prove that Σ^{im} generates $(\mathfrak{a}/\mathfrak{c}_a)^*$, where $\mathfrak{c}_a = \mathfrak{c} \cap \mathfrak{a}$ is the center of \mathfrak{m} . The natural homomorphism of vector spaces $\pi : \mathfrak{a} \rightarrow \mathfrak{h}/\mathfrak{c}$ induces a monomorphism $\bar{\pi} : \mathfrak{a}/\mathfrak{c}_a \rightarrow \mathfrak{h}/\mathfrak{c}$. By duality, the homomorphism $\bar{\pi}^* : (\mathfrak{h}/\mathfrak{c})^* \rightarrow (\mathfrak{a}/\mathfrak{c}_a)^*$ is surjective and $\bar{\pi}^*(\Delta^{im}) \subset \Sigma^{im}$ generates $(\mathfrak{a}/\mathfrak{c}_a)^*$.

3) Suppose that \mathfrak{g} is Lorentzian (hence of indefinite type) and let $(.,.)$ be an invariant non-degenerate bilinear form on \mathfrak{g} . Then, the restriction of $(.,.)$ to $\mathfrak{h}_{\mathbb{R}}$ has signature $(+ + \dots +, -)$ and any maximal totally isotropic subspace of $\mathfrak{h}_{\mathbb{R}}$ relatively to $(.,.)$ is one dimensional. Let $\mathfrak{a}_{\mathbb{R}} := \mathfrak{a} \cap \mathfrak{h}_{\mathbb{R}}$ and let $(.,.)_a$ be the restriction of $(.,.)$ to \mathfrak{m} . As \mathfrak{m} is of indefinite type, $\dim(\mathfrak{a}) \geq 2$ and the restriction of $(.,.)_a$ to $\mathfrak{a}_{\mathbb{R}}$ is non-null. It follows that the orthogonal subspace \mathfrak{m}^\perp

of \mathfrak{m} relatively to $(\cdot, \cdot)_a$ is a proper ideal of \mathfrak{m} . Since \mathfrak{m} is perfect (because \mathfrak{g} is) we deduce that $\mathfrak{m}^\perp = \{0\}$ (cf. [12, 1.7]) and the invariant bilinear form $(\cdot, \cdot)_a$ is non-degenerate. It follows that \mathfrak{m} is symmetrizable and the bilinear form $(\cdot, \cdot)_a$ when restricted to $\mathfrak{a}_\mathbb{R}$ is non-degenerate; since \mathfrak{m} is of indefinite type, it can not be positive definite. Hence, the bilinear form $(\cdot, \cdot)_a$ has signature $(+ + \dots +, -)$ on $\mathfrak{a}_\mathbb{R}$ and then the grading subalgebra \mathfrak{m} is Lorentzian. ■

Definition 3.7. Let Π_a be a root basis of Σ and let Σ^+ be the corresponding set of positive roots. The root basis is said to be adapted to the root basis Π of Δ if $\rho_a(\Delta^+) \subset \Sigma^+ \cup \{0\}$.

We will see (3.10) that adapted root bases always exist.

Lemma 3.8. Let Π_a be a root basis of Σ such that $\rho_a(\Delta_+^{im}) \subset \Sigma_+^{im}$ and let X_a be the corresponding positive Tits cone. Then we have $\bar{X}_a \subset \bar{X} \cap \mathfrak{a}$.

Proof. As $\Delta^{im} \neq \emptyset$, one has $\bar{X} = \{h \in \mathfrak{h}_\mathbb{R}; \langle \alpha, h \rangle \geq 0, \forall \alpha \in \Delta_+^{im}\}$ (see Remark 1.3). The lemma follows from Lemma 3.4. ■

Lemma 3.9. Suppose that $\Delta^{im} \neq \emptyset$. Let $p \in \bar{X}$ such that $\langle \alpha, p \rangle \in \mathbb{Z}, \forall \alpha \in \Delta$, and $\langle \beta, p \rangle > 0, \forall \beta \in \Delta_+^{im}$. Then $p \in \overset{\circ}{X}$.

Proof. The result is clear when Δ is of affine type since

$$\overset{\circ}{X} = \bar{X} = \{h \in \mathfrak{h}_\mathbb{R}; \langle \delta, h \rangle > 0\}.$$

Suppose now that Δ is of indefinite type. If one looks to the proof of Proposition 5.8.c) in [12], one can show that an element $p \in \bar{X}$ satisfying the conditions of the lemma lies in X . As Δ_+^{im} is W -invariant, we may assume that p lies in the fundamental chamber C . Hence there exists a subset J of I such that $\{\alpha \in \Delta; \langle \alpha, p \rangle = 0\} = \Delta_J = \Delta \cap \sum_{j \in J} \mathbb{Z}\alpha_j$. Since $\Delta_J \cap \Delta^{im} = \emptyset$, the root subsystem Δ_J is of finite type and p lies in the finite type facet of type J . Thus $p \in \overset{\circ}{X}$ (see section 1). ■

Theorem 3.10. There exists a root basis Π_a of Σ which is adapted to the root basis Π of Δ . Moreover, there exists a finite type subset J of I such that $\Delta_J = \{\alpha \in \Delta; \rho_a(\alpha) = 0\}$.

N.B. This is part 1) of Theorem 2.

Proof. Let $\Pi_a = \{\gamma_i, i \in \bar{I}\}$ be a root basis of Σ such that $\rho_a(\Delta_+^{im}) \subset \Sigma_+^{im}$, where \bar{I} is just a set indexing the basis elements. Let $p \in \mathfrak{a}$ such that $\langle \gamma_i, p \rangle = 1, \forall i \in \bar{I}$ and let $P = \{\alpha \in \Delta; \langle \alpha, p \rangle \geq 0\}$. If Δ is finite, then P is clearly a parabolic subsystem of Δ and the result is trivial. Suppose now that $\Delta^{im} \neq \emptyset$; then p satisfies the conditions of the Lemma 3.9 and we may assume that p lies in the facet of type J for some subset J of finite type in I . In which case $P = \Delta_J \cup \Delta^+$ is the standard parabolic subsystem of finite type J . Note that, for

$\gamma \in \Sigma^+$, one has $\langle \gamma, p \rangle = ht_a(\gamma)$ the height of γ with respect to Π_a . It follows that $\{\alpha \in \Delta; \rho_a(\alpha) = 0\} = \Delta_J$, in particular, $\rho_a(\Delta^+) = \rho_a(P) \subset \Sigma^+ \cup \{0\}$. Hence, the root basis Π_a is adapted to Π . ■

Corollary 3.11. Σ is indecomposable.

Proof. For $\gamma_1, \gamma_2 \in \Pi_a$, there are $\alpha_1, \alpha_2 \in \Delta_+$ such that $\gamma_i = \rho_a(\alpha_i)$. But γ_i is not a sum in Σ_+ , so, up to Δ_J , α_i is not a sum: we may assume $\alpha_i \in \Pi$. As Δ is indecomposable, there is a root $\alpha \in \Delta \cap (\alpha_1 + \alpha_2 + \sum_{\alpha \in \Pi} \mathbb{Z}^+ \alpha)$. Now $\rho_a(\alpha) \in (\Sigma \cup \{0\}) \cap (\gamma_1 + \gamma_2 + \sum_{\gamma \in \Pi_a} \mathbb{Z}^+ \gamma) \subset \Sigma$ and γ_1, γ_2 have to be in the same connected component of Π_a . ■

From now on, we fix a root basis $\Pi_a = \{\gamma_s, s \in \bar{I}\}$, for the grading root system Σ , which is adapted to the root basis $\Pi = \{\alpha_i, i \in I\}$ of Δ (see Theorem 3.10). As before, let $J := \{j \in I; \rho_a(\alpha_j) = 0\}$ and $I' := I \setminus J$. For $k \in I'$, we denote, as above, by I_k the connected component of $J \cup \{k\}$ containing k , and $J_k := J \cap I_k$.

Proposition 3.12.

- 1) Let $s \in \bar{I}$, then there exists $k_s \in I'$ such that $\rho_a(\alpha_{k_s}) = \gamma_s$ and any preimage $\alpha \in \Delta$ of γ_s is equal to α_{k_s} modulo $\sum_{j \in J_k} \mathbb{Z} \alpha_j$ for some $k \in I'$ satisfying $\rho_a(\alpha_k) = \gamma_s$.
- 2) Let $k \in I'$ such that $\rho_a(\alpha_k)$ is a real root of Σ . Then $\rho_a(\alpha_k) \in \Pi_a$ is a simple root.

Proof. This result was proved by J. Nervi for affine algebras (see [17, 2.3.10] and the proof of Prop. 2.3.12). The arguments used there are available for general Kac-Moody algebras. ■

We introduce the following notations :

$$I'_{re} := \{i \in I'; \rho_a(\alpha_i) \in \Pi_a\}; \quad I'_{im} := I' \setminus I'_{re},$$

$$I_{re} = \bigcup_{k \in I'_{re}} I_k; \quad J_{re} = I_{re} \cap J = \bigcup_{k \in I'_{re}} J_k; \quad J^\circ = J \setminus J_{re}$$

$$\Gamma_s := \{i \in I'; \rho_a(\alpha_i) = \gamma_s\}, \forall s \in \bar{I}.$$

Note that J° is not connected to I_{re} .

Remark 3.13.

- 1) In view of Proposition 3.12, assertion 2), one has $\rho_a(\alpha_k) \in \Sigma_+^{im}, \forall k \in I'_{im}$.
- 2) $I = I_{re} \cup I'_{im} \cup J^\circ$ is a disjoint union.
- 3) If $I'_{im} = \emptyset$, then $I = I_{re} \cup J^\circ$. Since I is connected (and I_{re} is not connected to J°) we deduce that $J^\circ = \emptyset, I = I_{re}$ and $I'_{re} = I' = I \setminus J$.
- 4) If $I'_{im} \neq \emptyset$, then I_{re} may be non-connected (see the example in §5 below).

Proposition 3.14.

- 1) Let $k \in I'_{re}$, then I_k is of finite type.

- 2) Let $s \in \bar{I}$. If $|\Gamma_s| \geq 2$ and $k \neq l \in \Gamma_s$, then $I_k \cup I_l$ is not connected: $\mathfrak{g}(I_k)$ and $\mathfrak{g}(I_l)$ commute and are orthogonal.
- 3) For all $k \in I'_{re}$, (I_k, J_k) is an irreducible C -admissible pair.
- 4) The derived subalgebra \mathfrak{m}' of the grading algebra \mathfrak{m} is contained in $\mathfrak{g}'(I_{re})$ (as defined in proposition 1.2).

Proof.

- 1) Suppose that there exists $k \in I'_{re}$ such that I_k is not of finite type; then there exists an imaginary root β_k whose support is the whole I_k . Hence, there exists a positive integer $m_k \in \mathbb{N}$ such that $\rho_a(\beta_k) = m_k \rho(\alpha_k)$ is an imaginary root of Σ . It follows that $\rho_a(\alpha_k)$ is an imaginary root and this contradicts the fact that $k \in I'_{re}$.
- 2) Let $s \in \bar{I}$ such that $|\Gamma_s| \geq 2$ and let $k \neq l \in \Gamma_s$. Since $V_{n\gamma_s} = \{0\}$ for all integer $n \geq 2$, the same argument used in 1) shows that $I_k \cup I_l$ is not connected, and I_k and I_l are its two connected components. In particular, $[\mathfrak{g}(I_k), \mathfrak{g}(I_l)] = \{0\}$ and $(\mathfrak{g}(I_k), \mathfrak{g}(I_l)) = \{0\}$.
- 3) Let $k \in I'_{re}$ and let $s \in \bar{I}$ such that $\rho_a(\alpha_k) = \gamma_s$. Let $(\bar{X}_s, \bar{H}_s = \gamma_s^\check, \bar{Y}_s)$ be an \mathfrak{sl}_2 -triple in \mathfrak{m} corresponding to the simple root γ_s . Let V_{γ_s} be the weight space of \mathfrak{g} corresponding to γ_s . In view of Proposition 3.12, assertion 1), one has :

$$V_{\gamma_s} = \bigoplus_{l \in \Gamma_s} V_{\gamma_s} \cap \mathfrak{g}(I_l). \tag{3.1}$$

Hence, one can write :

$$\bar{X}_s = \sum_{l \in \Gamma_s} E_l; \quad \bar{Y}_s = \sum_{l \in \Gamma_s} F_l, \tag{3.2}$$

with $E_l \in V_{\gamma_s} \cap \mathfrak{g}(I_l)$ and $F_l \in V_{-\gamma_s} \cap \mathfrak{g}(I_l)$. It follows from assertion 2) that

$$\bar{H}_s = \gamma_s^\check = [\bar{X}_s, \bar{Y}_s] = \sum_{l \in \Gamma_s} [E_l, F_l] = \sum_{l \in \Gamma_s} H_l, \tag{3.3}$$

where $H_l := [E_l, F_l] \in \mathfrak{h}_{I_l}, \forall l \in \Gamma_s$. Then one has, for $k \in \Gamma_s$,

$$2 = \langle \gamma_s, \gamma_s^\check \rangle = \langle \alpha_k, \gamma_s^\check \rangle = \sum_{l \in \Gamma_s} \langle \alpha_k, H_l \rangle = \langle \alpha_k, H_k \rangle,$$

and for $j \in J_k$,

$$0 = \langle \alpha_j, \gamma_s^\check \rangle = \sum_{l \in \Gamma_s} \langle \alpha_j, H_l \rangle = \langle \alpha_j, H_k \rangle.$$

In particular, H_k is the unique semi-simple element of \mathfrak{h}_{I_k} satisfying :

$$\langle \alpha_i, H_k \rangle = 2\delta_{i,k}, \forall i \in I_k. \tag{3.4}$$

Hence, (E_k, H_k, F_k) is an \mathfrak{sl}_2 -triple in the simple Lie algebra $\mathfrak{g}(I_k)$ and since $V_{2\gamma_s} = \{0\}$, (I_k, J_k) is an irreducible C -admissible pair for all $k \in \Gamma_s$. The statement 4) follows from the relation (3.2). ■

Corollary 3.15. *The pair (I_{re}, J_{re}) is C -admissible (in the eventually decomposable sense of Remark 2.15). If $I'_{im} = \emptyset$, then $I_{re} = I$, $J_{re} = J$ and \mathfrak{g} is finitely Δ^J -graded, with grading subalgebra \mathfrak{g}^J .*

N.B. We have got part 2) of Theorem 2.

Proof. The first assertion is a consequence of Proposition 3.14. By Remark 3.13, when $I'_{im} = \emptyset$, we have $I = I_{re}$; hence, by Theorem 2.14, \mathfrak{g} is finitely Δ^J -graded. ■

Definition 3.16. If $I'_{im} \neq \emptyset$, then (I, J) is called a generalized C -admissible pair and the gradation of \mathfrak{g} by Σ and \mathfrak{m} is said imaginary.

On the contrary if $I'_{im} = \emptyset$, the gradation is said real.

If $I'_{im} = J = \emptyset$, the Kac-Moody algebra \mathfrak{g} is said to be maximally finitely Σ -graded.

Corollary 3.17. *The grading subalgebra \mathfrak{m} of \mathfrak{g} is symmetrizable and the restriction to \mathfrak{m} of the invariant bilinear form of \mathfrak{g} is non-degenerate.*

Proof. Let $(\cdot, \cdot)_a$ be the restriction to \mathfrak{m} of the invariant bilinear form (\cdot, \cdot) of \mathfrak{g} . Recall from the proof of Proposition 3.14 that $\gamma_s = \sum_{k \in \Gamma_s} H_k, \forall s \in \bar{I}$. In particular $(\gamma_s, \gamma_s)_a = \sum_{k \in \Gamma_s} (H_k, H_k) > 0$. It follows that $(\cdot, \cdot)_a$ is a non-degenerate invariant bilinear form on \mathfrak{m} (see §1) and that \mathfrak{m} is symmetrizable. ■

Corollary 3.18. *Let \mathfrak{h}^J be the orthogonal of \mathfrak{h}_J in \mathfrak{h} . For $k \in I'_{im}$, write*

$$\rho_a(\alpha_k) = \sum_{s \in \bar{I}} n_{s,k} \gamma_s.$$

For $s \in \bar{I}$, choose l_s a representative element of Γ_s . Then $\mathfrak{a}/\mathfrak{c}_a$ can be viewed as the subspace of $\mathfrak{h}^J/\mathfrak{c}$ defined by the following relations :

$$\langle \alpha_k, h \rangle = \langle \alpha_{l_s}, h \rangle, \forall k \in \Gamma_s, \forall s \in \bar{I}$$

$$\langle \alpha_k, h \rangle = \sum_{s \in \bar{I}} n_{s,k} \langle \alpha_{l_s}, h \rangle, \forall k \in I'_{im}.$$

Proof. The subspace of $\mathfrak{h}^J/\mathfrak{c}$ defined by the above relations has dimension $|\bar{I}|$ and contains $\mathfrak{a}/\mathfrak{c}_a$ and hence it is equal to $\mathfrak{a}/\mathfrak{c}_a$. ■

Proposition 3.19. *Let $(\cdot, \cdot)_a$ be the restriction to \mathfrak{m} of the invariant bilinear form (\cdot, \cdot) of \mathfrak{g} .*

1) *Let $\mathfrak{a}' = \mathfrak{a} \cap \mathfrak{m}'$ and let \mathfrak{a}'' be a supplementary subspace of \mathfrak{a}' in \mathfrak{a} which is totally isotropic relatively to $(\cdot, \cdot)_a$. Then $\mathfrak{a}'' \cap \mathfrak{h}' = \{0\}$.*

2) *Let $A_{I_{re}}$ be the submatrix of A indexed by I_{re} . Then there exists a subspace $\mathfrak{h}_{I_{re}}$ of \mathfrak{h} containing \mathfrak{a} such that $(\mathfrak{h}_{I_{re}}, \Pi_{I_{re}}, \Pi_{I_{re}})$ is a realization of $A_{I_{re}}$. In particular, the Kac-Moody subalgebra $\mathfrak{g}(I_{re})$ associated to this realization (in 1.2) contains the grading subalgebra \mathfrak{m} .*

- 3) The Kac-Moody algebra $\mathfrak{g}(I_{re})$ is finitely $\Delta(I_{re})^{J_{re}}$ -graded and its grading subalgebra is the subalgebra $\mathfrak{g}(I_{re})^{J_{re}}$ associated to the C -admissible pair (I_{re}, J_{re}) as in Proposition 2.11.
- 4) The Kac-Moody algebra $\mathfrak{g}(I_{re})^{J_{re}}$ contains \mathfrak{m} .

Proof.

- 1) Recall that the center \mathfrak{c}_a of \mathfrak{m} is contained in the center \mathfrak{c} of \mathfrak{g} . Since $\mathfrak{h}' = \mathfrak{c}^\perp$ and \mathfrak{c}_a is in duality with \mathfrak{a}'' relatively to $(\cdot, \cdot)_a$, we deduce that $\mathfrak{a}'' \cap \mathfrak{h}' = \{0\}$.
- 2) From the proofs of 3.17 and 3.14 we get $\gamma_s^\vee = \sum_{k \in \Gamma_s} H_k \in \sum_{k \in \Gamma_s} \mathfrak{h}_{I_k} = \mathfrak{h}'_{I_{re}}$. So $\mathfrak{c}_a \subset \mathfrak{a}' \subset \mathfrak{h}'_{I_{re}} \subset \mathfrak{h}'$. It follows that $(\mathfrak{h}'_{I_{re}} + \mathfrak{h}^{I_{re}})$ is contained in \mathfrak{c}_a^\perp the orthogonal subspace of \mathfrak{c}_a in \mathfrak{h} . Since $\mathfrak{a}'' \cap \mathfrak{c}_a^\perp = \{0\}$, one can choose a supplementary subspace $\mathfrak{h}''_{I_{re}}$ of $(\mathfrak{h}'_{I_{re}} + \mathfrak{h}^{I_{re}})$ containing \mathfrak{a}'' . Let $\mathfrak{h}_{I_{re}} = \mathfrak{h}'_{I_{re}} \oplus \mathfrak{h}''_{I_{re}}$, then, by Proposition 1.2, $(\mathfrak{h}_{I_{re}}, \Pi_{I_{re}}, \check{\Pi}_{I_{re}})$ is a realization of $A_{I_{re}}$.
- 3) As in Corollary 3.15, assertion 3) is a simple consequence of Theorem 2.14.
- 4) The algebra \mathfrak{a} is in $\mathfrak{h}_{I_{re}} \cap \Pi_J^\perp = (\mathfrak{h}_{I_{re}})^{J_{re}}$. By the proof of Proposition 3.14, for $s \in \bar{I}$, \bar{X}_s and \bar{Y}_s are linear combinations of the elements in $\{E_k, F_k \mid k \in \Gamma_s\} \subset \mathfrak{g}(I_{re})^{J_{re}}$. Hence $\mathfrak{g}(I_{re})^{J_{re}}$ contains all generators of \mathfrak{m} . ■

Lemma 3.20. *Let \mathfrak{l} be a Kac-Moody subalgebra of \mathfrak{g} containing \mathfrak{m} . Then \mathfrak{l} is finitely Σ -graded. In particular, the Kac-Moody subalgebra $\mathfrak{g}(I_{re})$ or $\mathfrak{g}(I_{re})^{J_{re}}$ is finitely Σ -graded.*

N.B. Proposition 3.19 and Lemma 3.20 finish the proof of Theorem 2.

Proof. Recall that the Cartan subalgebra \mathfrak{a} of \mathfrak{m} is $\text{ad}_{\mathfrak{g}}$ -diagonalizable. Since \mathfrak{l} is $\text{ad}(\mathfrak{a})$ -invariant, one has $\mathfrak{l} = \sum_{\gamma \in \Sigma \cup \{0\}} V_\gamma \cap \mathfrak{l}$. By assumption $\{0\} \neq \mathfrak{m}_\gamma \subset V_\gamma \cap \mathfrak{l}$ for all $\gamma \in \Sigma$. Thus, \mathfrak{l} is finitely Σ -graded. ■

Proposition 3.21. *If $I'_{im} = \emptyset$, then $\mathfrak{g}(I_{re}) = \mathfrak{g}$ and the C -admissible subalgebra \mathfrak{g}^J is maximally finitely Σ -graded, with grading subalgebra \mathfrak{m} .*

Proof. This result is due to J. Nervi ([17, 2.5.10]) for the affine case; it follows from the facts that $V_0 \cap \mathfrak{g}^J = \mathfrak{h}^J$ and $\mathfrak{m} \subset \mathfrak{g}^J$ (see Prop. 3.19). ■

We now want a precise description of the gradation of $\mathfrak{g}(I_{re})$ by Σ and \mathfrak{m} ; particularly in the case (already mentioned in Remark 3.13) where $\mathfrak{g}(I_{re})$ (and so $\mathfrak{g}(I_{re})^{J_{re}}$) is decomposable.

Let $I_{re}^1, I_{re}^2, \dots, I_{re}^q$ be the connected components of I_{re} and $J_{re}^i := J_{re} \cap I_{re}^i$, $i = 1, 2, \dots, q$. Then $\mathfrak{g}(I_{re}) = \bigoplus_{i=1}^q \mathfrak{g}(I_{re}^i)$ and hence $\mathfrak{g}(I_{re})^{J_{re}} = \bigoplus_{i=1}^q \mathfrak{g}(I_{re}^i)^{J_{re}^i}$ (see Remark 2.15).

Retain the notations introduced just before Proposition 3.14 and those introduced in its proof. For $s \in \bar{I}$ and $i = 1, 2, \dots, q$, let $\Gamma_s^i := \Gamma_s \cap I_{re}^i$. If Γ_s^i is non-empty, put $E_s^i := \sum_{l \in \Gamma_s^i} E_l$; $F_s^i := \sum_{l \in \Gamma_s^i} F_l$ and $H_s^i := \sum_{l \in \Gamma_s^i} H_l$. We take $E_s^i = F_s^i = H_s^i = 0$ if Γ_s^i is empty.

Note that $\Gamma_s = \bigcup_{i=1}^q \Gamma_s^i$ (disjoint union) and from the proof of the Proposition 3.14 we get the following relations :

$$\bar{X}_s = \sum_{i=1}^q E_s^i; \quad \bar{Y}_s = \sum_{i=1}^q F_s^i, \forall s \in \bar{I}, \tag{3.5}$$

$$\bar{H}_s = \gamma_{\check{s}} = [\bar{X}_s, \bar{Y}_s] = \sum_{i=1}^q [E_s^i, F_s^i] = \sum_{i=1}^q H_s^i, \forall s \in \bar{I}. \tag{3.6}$$

Lemma 3.22. *Let $s \in \bar{I}$ and $i \in \{1, 2, \dots, q\}$ such that $\Gamma_s^i \neq \emptyset$. Then we have*
 1) $\Gamma_t^i \neq \emptyset$ for all $t \in \bar{I}$ satisfying $\langle \gamma_t, \gamma_{\check{s}} \rangle < 0$.
 2) $\Gamma_t^i \neq \emptyset, \forall t \in \bar{I}$.

Proof. To prove 1), suppose $\Gamma_t^i = \emptyset$ for any t satisfying $\langle \gamma_t, \gamma_{\check{s}} \rangle < 0$. Let $k \in \Gamma_s^i$, then $\langle \gamma_s, \gamma_{\check{t}} \rangle = \sum_{\substack{j=1 \\ j \neq i}}^q \langle \alpha_k, H_t^j \rangle = 0$, a contradiction since $\langle \gamma_s, \gamma_{\check{t}} \rangle$ must

be negative. Thus $\Gamma_s^i \neq \emptyset$ iff $\Gamma_t^i \neq \emptyset$. The second statement follows from the connectedness of \bar{I} : For $t \in \bar{I}$, there exists a sequence $s_0 = s, s_1, \dots, s_n = t$ in \bar{I} such that s_j is linked to s_{j+1} for all $j = 0, 1, \dots, n - 1$. By 1) $\Gamma_{s_j}^i$ is, as Γ_s^i , non-empty for all $j = 0, 1, \dots, n$. In particular $\Gamma_t^i \neq \emptyset$. ■

Lemma 3.23. $\Gamma_s^i \neq \emptyset, \forall s \in \bar{I}$, and $(H_s^i)_{s \in \bar{I}}$ is free for all $i = 1, 2, \dots, q$.

Proof. Recall that $I_{re} = \bigcup_{k \in I'_{re}} I_k$, with all the I_k connected. Let $i \in \{1, 2, \dots, q\}$ and let $k \in I'_{re}$ such that $I_k \subset I_{re}^i$. Let $s \in \bar{I}$ such that $\rho_a(\alpha_k) = \gamma_s$, then $k \in \Gamma_s^i$ and $\Gamma_s^i \neq \emptyset$. By the Lemma 3.22, $\Gamma_t^i \neq \emptyset$ for all $t \in \bar{I}$. Thus $H_s^i \neq 0, \forall s \in \bar{I}; \forall i = 1, 2, \dots, q$, and the freeness of $(H_s^i)_{s \in \bar{I}}$ follows from that of $(H_k)_{k \in I'_{re}}$. ■

Proposition 3.24. *For $i = 1, 2, \dots, q$, let p_i be the projection of $\mathfrak{g}(I_{re})$ on $\mathfrak{g}(I_{re}^i)$ with kernel $\bigoplus_{j \neq i} \mathfrak{g}(I_{re}^j)$ and let $\mathfrak{m}_i := p_i(\mathfrak{m})$. Then we have :*

- 1) \mathfrak{m}_i is a Kac-Moody subalgebra of $\mathfrak{g}(I_{re}^i)^{J_{re}^i}$ isomorphic to \mathfrak{m} .
- 2) The Kac-Moody subalgebra $\mathfrak{g}(I_{re}^i)^{J_{re}^i}$ is maximally finitely Σ_i -graded, where Σ_i is the root system of \mathfrak{m}_i relative to the Cartan subalgebra $\mathfrak{a}_i := p_i(\mathfrak{a})$.

N.B. Note that \mathfrak{m} is contained in $\bigoplus_{i=1}^q \mathfrak{m}_i$. In particular, $\bigoplus_{i=1}^q \mathfrak{m}_i$ is finitely Σ -graded.

If we identify $\bigoplus_{i=1}^q \mathfrak{m}_i$ with \mathfrak{m}^q , then the grading subalgebra \mathfrak{m} can be viewed as the diagonal subalgebra $\Delta(\mathfrak{m}^q)$ of \mathfrak{m}^q : $\Delta(\mathfrak{m}^q) := \{(X, X, \dots, X) ; X \in \mathfrak{m}\}$.

Proof. For $i \in \{1, 2, \dots, q\}$, p_i is a morphism of Lie algebras and $\mathfrak{m}_i := p_i(\mathfrak{m})$ is contained in $\mathfrak{g}(I_{re}^i)^{J_{re}^i}$. For $s \in \bar{I}$, one has $p_i(\gamma_{\check{s}}) = H_s^i$. Thus the restriction of p_i to $\mathfrak{a}' := [\mathfrak{a}, \mathfrak{a}] = \bigoplus_{s \in \bar{I}} \mathbb{C}\gamma_{\check{s}}$ is injective by Lemma 3.23. Since \mathfrak{m} is indecomposable, p_i when restricted to \mathfrak{m} is still injective (see [12, 1.7]). Thus $\mathfrak{m}_i = p_i(\mathfrak{m})$ is isomorphic to \mathfrak{m} and we have the following commutative diagram :

$$\begin{array}{ccc}
 \mathfrak{m} & \xrightarrow[\sim]{p_i} & \mathfrak{m}_i \\
 \downarrow & \circlearrowleft & \downarrow \\
 \mathfrak{g}(I_{re})^{J_{re}} & \xrightarrow{p_i} & \mathfrak{g}(I_{re}^i)^{J_{re}^i} \\
 \downarrow & \circlearrowleft & \downarrow \\
 \mathfrak{g}(I_{re}) & \xrightarrow{p_i} & \mathfrak{g}(I_{re}^i)
 \end{array}$$

For the second assertion, Let $\mathfrak{a}_i := p_i(\mathfrak{a})$ and $\Sigma_i = \Delta(\mathfrak{m}_i, \mathfrak{a}_i)$. When restricted to \mathfrak{m} , p_i induces an isomorphism of root systems $\psi_i : \Sigma_i \rightarrow \Sigma$ such that

$$\langle \alpha, a \rangle = \langle \psi_i^{-1}(\alpha), p_i(a) \rangle, \quad \forall \alpha \in \Sigma, \forall a \in \mathfrak{a}.$$

Note that for $\alpha \in \Sigma$ and $X \in \mathfrak{g}(I_{re})$ satisfying $[a, X] = \langle \alpha, a \rangle X, \forall a \in \mathfrak{a}$, one has $[a_i, p_i(X)] = \langle \psi_i^{-1}(\alpha), a_i \rangle p_i(X), \forall a_i \in \mathfrak{a}_i$. Since p_i is surjective and $\mathfrak{g}(I_{re})$ (resp. $\mathfrak{g}(I_{re})^{J_{re}}$) is finitely Σ -graded, the Kac-Moody subalgebra $\mathfrak{g}(I_{re}^i)$ (resp. $\mathfrak{g}(I_{re}^i)^{J_{re}^i}$) is also finitely Σ_i -graded. For $k \in I_{re}^i$, Let $\rho_i(\alpha_k)$ be the restriction of α_k to \mathfrak{a}_i . Then $(\rho_i(\alpha_k) = 0) \iff (\rho_a(\alpha_k) = 0) \iff (k \in J_{re}^i)$. By Proposition 3.21, $\mathfrak{g}(I_{re}^i)^{J_{re}^i}$ is maximally finitely Σ_i -graded. ■

Corollary 3.25. *If \mathfrak{g} is Lorentzian then I_{re} is connected.*

Proof. If \mathfrak{g} is Lorentzian, then by Proposition 3.6, the grading subalgebra \mathfrak{m} and hence all the \mathfrak{m}_i ($i = 1, 2, \dots, q$) are also Lorentzian. When restricted to $\bigoplus_{i=1}^q \mathfrak{a}_i$, the invariant bilinear form (\cdot, \cdot) is still non-degenerate and has signature $(q(r - 1), q)$, where r is the common rank of the $\mathfrak{m}_i, i = 1, 2, \dots, q$. Hence $q = 1$ and I_{re} is connected. ■

Proposition 3.26. *If \mathfrak{g} is of finite, affine or hyperbolic type, then any finite gradation is real: $I'_{im} = \emptyset$ and (I, J) is a C -admissible pair.*

Proof. The result is trivial if \mathfrak{g} is of finite type. Suppose $I'_{im} \neq \emptyset$ for one of the other cases. If \mathfrak{g} is affine, then I_{re} is of finite type and by Lemma 3.19, \mathfrak{m} is contained in the finite dimensional semi-simple Lie algebra $\mathfrak{g}(I_{re})$. This contradicts the fact that \mathfrak{m} is, as \mathfrak{g} , of affine type (see Proposition 3.6). If \mathfrak{g} is hyperbolic, then it is Lorentzian and perfect (cf. section 1). By Lemma 3.20 and Corollary 3.25, $\mathfrak{g}(I_{re})$ is an indecomposable finitely Σ -graded Kac-Moody subalgebra of \mathfrak{g} . As I_{re} is assumed to be a proper connected subset of I , $\mathfrak{g}(I_{re})$ is of finite or affine type, a contradiction since, by Proposition 3.6, \mathfrak{m} must be Lorentzian. Hence $I'_{im} = \emptyset$ in the two last cases. ■

Proposition 3.27. *If \mathfrak{g} is hyperbolic, then the grading subalgebra \mathfrak{m} is also hyperbolic.*

Proof. Recall that in this case $I_{re} = I$ (see Proposition 3.26 and Corollary 3.15). Let \bar{I}^1 be a proper subset of \bar{I} and suppose that \bar{I}^1 is connected.

Let $I^1 = \bigcup_{s \in \bar{I}^1} (\bigcup_{k \in \Gamma_s} I_k)$. Then, I^1 is a proper subset of I . We may assume that the subalgebra $\mathfrak{m}(\bar{I}^1)$ of \mathfrak{m} is contained in $\mathfrak{g}(I^1)$. Let $\Sigma^1 := \Sigma(\bar{I}^1)$ be the root system of $\mathfrak{m}(\bar{I}^1)$. Then, it is not difficult to check that $\mathfrak{g}(I^1)$ is finitely Σ^1 -graded. The argument used in Proposition 3.24 shows that the indecomposable components of $\mathfrak{g}(I^1)$ (which all are of finite or affine type) are finitely Σ^1 -graded. By Proposition 3.6, $\mathfrak{m}(\bar{I}^1)$ is of finite or affine type. Hence \mathfrak{m} is hyperbolic. ■

Corollary 3.28. *The problem of classification of finite real gradations of \mathfrak{g} comes down first to classify the C -admissible pairs (I, J) of \mathfrak{g} and then the maximal finite gradations of the corresponding admissible algebra \mathfrak{g}^J . When \mathfrak{g} is of finite, affine or hyperbolic type, we get thus all finite gradations.*

Proof. This follows from Proposition 3.26, Proposition 3.21 and Lemma 1.5. ■

4. Maximal gradations

We assume now moreover that \mathfrak{g} is maximally finitely Σ -graded. We keep the notations in section 3 but we have $J = I'_{im} = \emptyset$. So \bar{I} is a quotient of I , with quotient map ρ defined by $\rho_a(\alpha_k) = \gamma_{\rho(k)}$. For $s \in \bar{I}$, $\Gamma_s = \rho^{-1}(\{s\})$.

Proposition 4.1.

- 1) If $k \neq l \in I$ and $\rho(k) = \rho(l)$, then there is no link between k and l in the Dynkin diagram of A : $\alpha_k(\alpha_l^\vee) = \alpha_l(\alpha_k^\vee) = 0$ and $(\alpha_k, \alpha_l) = 0$.
- 2) $\mathfrak{a} \subset \{h \in \mathfrak{h} \mid \alpha_k(h) = \alpha_l(h) \text{ whenever } \rho(k) = \rho(l)\}$.
- 3) For good choices of the simple coroots and Chevalley generators $(\alpha_k^\vee, e_k, f_k)_{k \in I}$ in \mathfrak{g} and $(\gamma_s^\vee, \bar{X}_s, \bar{Y}_s)_{s \in \bar{I}}$ in \mathfrak{m} , we have $\gamma_s^\vee = \sum_{k \in \Gamma_s} \alpha_k^\vee$, $\bar{X}_s = \sum_{k \in \Gamma_s} e_k$ and $\bar{Y}_s = \sum_{k \in \Gamma_s} f_k$.
- 4) In particular, for $s, t \in \bar{I}$, we have $\gamma_s(\gamma_t^\vee) = \sum_{k \in \Gamma_t} \alpha_i(\alpha_k^\vee)$ for any $i \in \Gamma_s$.

Proof. Assertions 1) and 2) are proved in 3.14 and 3.18. For $i \in \Gamma_s$, $\gamma_s = \rho_a(\alpha_i)$ is the restriction of α_i to \mathfrak{a} ; so 4) is a consequence of 3).

For 3) recall the proof of Proposition 3.14. The \mathfrak{sl}_2 -triple $(\bar{X}_s, \gamma_s^\vee, \bar{Y}_s)$ may be written $\gamma_s^\vee = \sum_{k \in \Gamma_s} H_k$, $\bar{X}_s = \sum_{k \in \Gamma_s} E_k$ and $\bar{Y}_s = \sum_{k \in \Gamma_s} F_k$ where (E_k, H_k, F_k) is an \mathfrak{sl}_2 -triple in $\mathfrak{g}(I_k)$, with $\alpha_k(H_k) = 2$. But now $J = I'_{im} = \emptyset$, so $I_k = \{k\}$ and $\mathfrak{g}(I_k) = \mathbb{C}e_k \oplus \mathbb{C}\alpha_k^\vee \oplus \mathbb{C}f_k$, hence the result. ■

So the grading subalgebra \mathfrak{m} may be entirely described by the quotient map ρ . We look now to the reciprocal construction.

So \mathfrak{g} is an indecomposable and symmetrizable Kac-Moody algebra associated to a generalized Cartan matrix $A = (a_{i,j})_{i,j \in I}$. We consider a quotient \bar{I} of I with quotient map $\rho : I \rightarrow \bar{I}$ and fibers $\Gamma_s = \rho^{-1}(\{s\})$ for $s \in \bar{I}$. We suppose that ρ is an admissible quotient i.e. that it satisfies the following two conditions :

(MG1) If $k \neq l \in I$ and $\rho(k) = \rho(l)$, then $a_{k,l} = \alpha_l(\alpha_k^\vee) = 0$.

(MG2) If $s \neq t \in \bar{I}$, then $\bar{a}_{s,t} := \sum_{i \in \Gamma_s} a_{i,j} = \sum_{i \in \Gamma_s} \alpha_j(\alpha_i^\vee)$ is independent of

the choice of $j \in \Gamma_t$.

Proposition 4.2. *The matrix $\bar{A} = (\bar{a}_{s,t})_{s,t \in \bar{I}}$ is an indecomposable generalized Cartan matrix.*

Proof. Let $s \neq t \in \bar{I}$ and let $j \in \Gamma_t$. By (MG1) one has $\bar{a}_{t,t} = \sum_{i \in \Gamma_t} a_{i,j} = a_{j,j} = 2$, and by (MG2) $\bar{a}_{s,t} := \sum_{i \in \Gamma_s} a_{i,j} \in \mathbb{Z}^-$ ($\forall j \in \Gamma_t$). Moreover, $\bar{a}_{s,t} = 0$ if and only if $a_{i,j} = 0$ ($= a_{j,i}$), $\forall (i,j) \in \Gamma_s \times \Gamma_t$. It follows that $\bar{a}_{s,t} = 0$ if and only if $\bar{a}_{t,s} = 0$, and \bar{A} is a generalized Cartan matrix. Since A is indecomposable, \bar{A} is also indecomposable. ■

Let $\mathfrak{h}^\Gamma = \{h \in \mathfrak{h} \mid \alpha_k(h) = \alpha_l(h) \text{ whenever } \rho(k) = \rho(l)\}$, $\gamma_s^\vee = \sum_{k \in \Gamma_s} \alpha_k^\vee$ and $\mathfrak{a}' = \bigoplus_{s \in \bar{I}} \mathbb{C}\gamma_s^\vee \subset \mathfrak{h}^\Gamma$. We may choose a subspace \mathfrak{a}'' in \mathfrak{h}^Γ such that $\mathfrak{a}'' \cap \mathfrak{a}' = \{0\}$, the restrictions $\bar{\alpha}_i =: \gamma_{\rho(i)}$ to $\mathfrak{a} = \mathfrak{a}' \oplus \mathfrak{a}''$ of the simple roots α_i (corresponding to different $\rho(i) \in \bar{I}$) are linearly independent and \mathfrak{a}'' is minimal for these two properties.

Proposition 4.3. *$(\mathfrak{a}, \{\gamma_s \mid s \in \bar{I}\}, \{\gamma_s^\vee \mid s \in \bar{I}\})$ is a realization of \bar{A} .*

Proof. Let ℓ be the rank of \bar{A} . Note that \mathfrak{a} contains $\mathfrak{a}' = \bigoplus_{s \in \bar{I}} \mathbb{C}\gamma_s^\vee$; the family $(\gamma_s)_{s \in \bar{I}}$ is free in the dual space \mathfrak{a}^* of \mathfrak{a} and satisfies $\langle \gamma_t, \gamma_s^\vee \rangle = \bar{a}_{s,t}$, $\forall s, t \in \bar{I}$. It follows that $\dim(\mathfrak{a}) \geq 2|\bar{I}| - \ell$ (see [11, 14.1] or [12, Ex. 1.3]). As \mathfrak{a} is minimal, we have $\dim(\mathfrak{a}) = 2|\bar{I}| - \ell$ (see [11, 14.2] for minimal realization). Hence $(\mathfrak{a}, \{\gamma_s \mid s \in \bar{I}\}, \{\gamma_s^\vee \mid s \in \bar{I}\})$ is a (minimal) realization of \bar{A} . ■

We note $\Delta^\rho = \Sigma \subset \bigoplus_{s \in \bar{I}} \mathbb{Z}\gamma_s$ the root system associated to this realization.

We define now $\bar{X}_s = \sum_{k \in \Gamma_s} e_k$ and $\bar{Y}_s = \sum_{k \in \Gamma_s} f_k$. Let $\mathfrak{m} = \mathfrak{g}^\rho$ be the Lie subalgebra of \mathfrak{g} generated by \mathfrak{a} and the elements \bar{X}_s, \bar{Y}_s for $s \in \bar{I}$.

Proposition 4.4. *The Lie subalgebra $\mathfrak{m} = \mathfrak{g}^\rho$ is the Kac-Moody algebra associated to the realization $(\mathfrak{a}, \{\gamma_s \mid s \in \bar{I}\}, \{\gamma_s^\vee \mid s \in \bar{I}\})$ of \bar{A} . Moreover, \mathfrak{g} is an integrable \mathfrak{g}^ρ -module with finite multiplicities.*

Proof. Clearly, the following relations hold in the Lie subalgebra \mathfrak{g}^ρ :

$$\begin{aligned} [\mathfrak{a}, \mathfrak{a}] &= 0, & [\bar{X}_s, \bar{Y}_t] &= \delta_{s,t} \gamma_s^\vee & (s, t \in \bar{I}); \\ [a, \bar{X}_s] &= \langle \gamma_s, a \rangle \bar{X}_s, & [a, \bar{Y}_s] &= -\langle \gamma_s, a \rangle \bar{Y}_s & (a \in \mathfrak{a}, s \in \bar{I}). \end{aligned}$$

For the Serre's relations, one has :

$$1 - \bar{a}_{s,t} \geq 1 - a_{i,j}, \quad \forall (i,j) \in \Gamma_s \times \Gamma_t.$$

In particular, one can see, by induction on $|\Gamma_s|$, that :

$$(\text{ad} \bar{X}_s)^{1-\bar{a}_{s,t}}(e_j) = \left(\sum_{i \in \Gamma_s} \text{ad} e_i \right)^{1-\bar{a}_{s,t}}(e_j) = 0, \quad \forall j \in \Gamma_t.$$

Hence

$$(\text{ad} \bar{X}_s)^{1-\bar{a}_{s,t}}(\bar{X}_t) = 0, \quad \forall s, t \in \bar{I},$$

and in the same way we obtain that :

$$(\text{ad } \bar{Y}_s)^{1-\bar{a}_{s,t}}(\bar{Y}_t) = 0, \forall s, t \in \bar{I}.$$

It follows that \mathfrak{g}^ρ is a quotient of the Kac-Moody algebra $\mathfrak{g}(\bar{A})$ associated to \bar{A} and $(\mathfrak{a}, \{\gamma_s \mid s \in \bar{I}\}, \{\gamma_s^\vee \mid s \in \bar{I}\})$ in which the Cartan subalgebra \mathfrak{a} of $\mathfrak{g}(\bar{A})$ is embedded. By [12, 1.7] \mathfrak{g}^ρ is equal to $\mathfrak{g}(\bar{A})$.

It's clear that \mathfrak{g} is an integrable \mathfrak{g}^ρ -module with finite dimensional weight spaces relative to the adjoint action of \mathfrak{a} , since for $\alpha = \sum_{i \in \bar{I}} n_i \alpha_i \in \Delta^+$, its restriction $\rho_a(\alpha)$ to \mathfrak{a} is given by

$$\rho_a(\alpha) = \sum_{s \in \bar{I}} \left(\sum_{i \in \Gamma_s} n_i \right) \gamma_s \tag{4.1}$$

■

Proposition 4.5. *The Kac-Moody algebra \mathfrak{g} is maximally finitely Δ^ρ -graded with grading subalgebra \mathfrak{g}^ρ .*

Proof. As in Theorem 2.14, we will see that $\rho_a(\Delta^+) \subset Q_+^\Gamma := \bigoplus_{s \in \bar{I}} \mathbb{Z}^+ \gamma_s$ satisfies,

as $\Sigma^+ = \Delta_+^\rho$, the following conditions :

- (i) $\gamma_s \in \rho_a(\Delta^+) \subset Q_+^\Gamma$, $2\gamma_s \notin \rho_a(\Delta^+)$, $\forall s \in \bar{I}$.
- (ii) if $\gamma \in \rho_a(\Delta^+)$, $\gamma \neq \gamma_s$, then the set $\{\gamma + k\gamma_s; k \in \mathbb{Z}\} \cap \rho_a(\Delta^+)$ is a string $\{\gamma - p\gamma_s, \dots, \gamma + q\gamma_s\}$, where $p, q \in \mathbb{Z}^+$ and $p - q = \langle \gamma, \gamma_s^\vee \rangle$;
- (iii) if $\gamma \in \rho_a(\Delta^+)$, then $\text{supp}(\gamma)$ is connected.

Clearly $\{\gamma_s \mid s \in \bar{I}\} \subset \rho_a(\Delta_+) \subset Q_+^\Gamma$. For $\alpha \in \Delta$ and $s \in \bar{I}$, the condition $\rho_a(\alpha) \in \mathbb{N}\gamma_s$ implies $\alpha \in \Delta(\Gamma_s)^+ = \{\alpha_i; i \in \Gamma_s\}$ [see (4.1)]. It follows that $2\gamma_s \notin \rho_a(\Delta_+)$ and (i) is satisfied. By Proposition 4.4, \mathfrak{g} is an integrable \mathfrak{g}^ρ -module with finite multiplicities. Hence, the propriety (ii) follows from [12, 3.6]. Let $\alpha \in \Delta_+$ and let $s, t \in \text{supp}(\rho_a(\alpha))$. By (4.1) there exists $(k, l) \in \Gamma_s \times \Gamma_t$ such that $k, l \in \text{supp}(\alpha)$, which is connected. Hence there exist $i_0 = k, i_1, \dots, i_{n+1} = l$ such that $\alpha_{i_j} \in \text{supp}(\alpha)$, $j = 0, 1, \dots, n+1$, and for $j = 0, 1, \dots, n$, i_j and i_{j+1} are linked relative to the generalized Cartan matrix A . In particular, $\rho(i_j) \neq \rho(i_{j+1}) \in \text{supp}(\rho_a(\alpha))$ and they are linked relative to the generalized Cartan matrix \bar{A} , $j = 0, 1, \dots, n$, with $\rho(i_0) = s$ and $\rho(i_{n+1}) = t$. Hence the connectedness of $\text{supp}(\rho_a(\alpha))$ relative to \bar{A} . It follows that $\rho_a(\Delta^+) = \Delta_+^\rho$ and hence $\rho_a(\Delta) = \Delta^\rho$ (see [12, Ex. 5.4]. In particular, \mathfrak{g} is finitely Δ^ρ -graded with $J = \emptyset = I'_{im}$.

■

Corollary 4.6. *The restriction to $\mathfrak{m} = \mathfrak{g}^\rho$ of the invariant bilinear form (\cdot, \cdot) of \mathfrak{g} is non-degenerate. In particular, the generalized Cartan matrix \bar{A} is symmetrizable of the same type as A .*

Proof. The first part of the corollary follows from Proposition 4.5 and Corollary 3.17. The second part follows from Proposition 3.6. ■

Remark 4.7. The map ρ coincides with the map (also denoted ρ) defined at the beginning of this section using the maximal gradation of Proposition 4.5. Conversely, Proposition 4.1 tells that, for a general maximal finite gradation, ρ is

admissible and $\mathfrak{m} = \mathfrak{g}^\rho$ for good choices of the Chevalley generators. So we get a good correspondence between maximal gradations and admissible quotient maps.

By Corollary 3.28 the real finite gradations of a Kac-Moody algebra \mathfrak{g} are bijectively associated to pairs of a C -admissible pair (I, J) and an admissible quotient map $\rho : I' = I \setminus J \rightarrow \bar{I}'$.

5. An example


The following example shows that imaginary gradations do exist. It shows in particular that, for a generalized C -admissible pair (I, J) , J° may be non-empty and I_{re} may be non-connected. Moreover, the Kac-Moody algebra \mathfrak{g} may be not graded by the root system of $\mathfrak{g}(I_{re})$.

The imaginary gradations will be studied in a forthcoming paper [7].


Example 5.1. Consider the Kac Moody algebra \mathfrak{g} corresponding to the indecomposable and symmetric generalized Cartan matrix A :

$$A = \begin{pmatrix} 2 & -3 & -1 & 0 & 0 & 0 \\ -3 & 2 & -1 & 0 & 0 & 0 \\ -1 & -1 & 2 & -1 & -1 & -1 \\ 0 & 0 & -1 & 2 & 0 & 0 \\ 0 & 0 & -1 & 0 & 2 & -3 \\ 0 & 0 & -1 & 0 & -3 & 2 \end{pmatrix}$$

with the corresponding Dynkin diagram :



Note that $\det(A) = 275$ and the symmetric submatrix of A indexed by $\{1, 2, 4, 5, 6\}$ has signature $(+++,-)$. Since $\det(A) > 0$, the matrix A should have signature $(++++,-)$. Let Σ be the root system associated to the strictly hyperbolic generalized Cartan matrix $\begin{pmatrix} 2 & -3 \\ -3 & 2 \end{pmatrix}$, the corresponding Dynkin diagram is the following :




We will see that \mathfrak{g} is finitely Σ -graded and describe the corresponding generalized C -admissible pair.

1) Let τ be the involutive diagram automorphism of \mathfrak{g} such that $\tau(1) = 5$, $\tau(2) = 6$ and τ fixes the other vertices. Let σ'_n be the normal semi-involution of \mathfrak{g} corresponding to the split real form of \mathfrak{g} . Consider the quasi-split real form $\mathfrak{g}_{\mathbb{R}}^1$ associated to the semi-involution $\tau\sigma'_n$ (see [2] or [6]). Then $\mathfrak{t}_{\mathbb{R}} := \mathfrak{h}_{\mathbb{R}}^{\tau}$ is a maximal split toral subalgebra of $\mathfrak{g}_{\mathbb{R}}^1$. The corresponding restricted root system $\Delta' := \Delta(\mathfrak{g}_{\mathbb{R}}, \mathfrak{t}_{\mathbb{R}})$ is reduced and the corresponding generalized Cartan matrix A' is

given by :

$$A' = \begin{pmatrix} 2 & -3 & -2 & 0 \\ -3 & 2 & -2 & 0 \\ -1 & -1 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{pmatrix}$$

with the corresponding Dynkin diagram :



Following N. Bardy [4, 9], there exists a split real Kac-Moody subalgebra $\mathfrak{m}_{\mathbb{R}}^1$ of $\mathfrak{g}_{\mathbb{R}}^1$ containing $\mathfrak{t}_{\mathbb{R}}$ such that $\Delta' = \Delta(\mathfrak{m}_{\mathbb{R}}^1, \mathfrak{t}_{\mathbb{R}})$. It follows that \mathfrak{g} is finitely Δ' -graded.

2) Let $\mathfrak{m}^1 := \mathfrak{m}_{\mathbb{R}}^1 \otimes \mathbb{C}$ and $\mathfrak{t} := \mathfrak{t}_{\mathbb{R}} \otimes \mathbb{C}$. Denote by $\alpha'_i := \alpha_i/\mathfrak{t}$, $i = 1, 2, 3, 4$. Put $\alpha'_1 = \alpha_1 + \alpha_5$, $\alpha'_2 = \alpha_2 + \alpha_6$, $\alpha'_3 = \alpha_3$ and $\alpha'_4 = \alpha_4$. Let $I^1 := \{1, 2, 3, 4\}$, then $(\mathfrak{t}, \Pi' = \{\alpha'_i, i \in I^1\}, \Pi^{\vee} = \{\alpha'_{\check{i}}, i \in I^1\})$ is a realization of A' which is symmetrizable and Lorentzian.

Let \mathfrak{m} be the Kac-Moody subalgebra of \mathfrak{m}^1 corresponding to the submatrix \bar{A} of A' indexed by $\{1, 2\}$. Thus $\bar{A} = \begin{pmatrix} 2 & -3 \\ -3 & 2 \end{pmatrix}$ is strictly hyperbolic. Let $\mathfrak{a} := \mathbb{C}\alpha'_1 \oplus \mathbb{C}\alpha'_2$ be the standard Cartan subalgebra of \mathfrak{m} and let $\Sigma = \Delta(\mathfrak{m}, \mathfrak{a})$. For $\alpha' \in \mathfrak{t}^*$, denote by $\rho_1(\alpha')$ the restriction of α' to \mathfrak{a} . Put $\gamma_s = \rho_1(\alpha'_s)$, $\gamma_{\check{s}} = \alpha'_{\check{s}}$, $s = 1, 2$. Then $\Pi_{\mathfrak{a}} = \{\gamma_1, \gamma_2\}$ is the standard root basis of Σ . One can see easily that $\rho_1(\alpha'_4) = 0$ and $\rho_1(\alpha'_3) = 2(\gamma_1 + \gamma_2)$ is a strictly positive imaginary root of Σ . Now we will show that \mathfrak{m}^1 is finitely Σ -graded.

Let $(\cdot, \cdot)_1$ be the normalized invariant bilinear form on \mathfrak{m}^1 such that short real roots have length 1 and long real roots have square length 2. Then there exists a positive rational q such that the restriction of $(\cdot, \cdot)_1$ to \mathfrak{t} has the matrix B_1 in the basis $\Pi^{\check{\cdot}}$, where :

$$B_1 = q \begin{pmatrix} 2 & -3 & -1 & 0 \\ -3 & 2 & -1 & 0 \\ -1 & -1 & 1 & -1/2 \\ 0 & 0 & -1/2 & 1 \end{pmatrix}$$

By duality, the restriction of $(\cdot, \cdot)_1$ to \mathfrak{t} induces a non-degenerate symmetric bilinear form on \mathfrak{t}^* (see [12, 2.1]) such that its matrix B'_1 in the basis Π' , is the following :

$$B'_1 = q^{-1} \begin{pmatrix} 2 & -3 & -2 & 0 \\ -3 & 2 & -2 & 0 \\ -2 & -2 & 4 & -2 \\ 0 & 0 & -2 & 4 \end{pmatrix}$$

Hence, q equals 2.

Note that for $\alpha' = \sum_{i=1}^4 n_i \alpha'_i \in \Delta'^+$, we have that

$$(\alpha', \alpha')_1 = n_1^2 + n_2^2 + 2n_3^2 + 2n_4^2 - 3n_1n_2 - 2n_1n_3 - 2n_2n_3 - 2n_3n_4. \tag{5.1}$$

We will show that $\rho_1(\Delta'^+) = \Sigma^+ \cup \{0\}$. Note that Σ can be identified with $\Delta' \cap (\mathbb{Z}\alpha'_1 + \mathbb{Z}\alpha'_2)$; hence ρ_1 is injective on Σ and $\Sigma^+ \subset \rho_1(\Delta'^+)$.

Let $(\cdot, \cdot)_a$ be the normalized invariant bilinear form on \mathfrak{m} such that all real roots have length 2. Then the restriction of $(\cdot, \cdot)_a$ to \mathfrak{a} has the matrix B_a in the basis $\Pi_a = \{\gamma_1, \gamma_2\}$, where :

$$B_a = \begin{pmatrix} 2 & -3 \\ -3 & 2 \end{pmatrix}$$

Since \bar{A} is symmetric, the non-degenerate symmetric bilinear form, on \mathfrak{a}^* , induced by the restriction of $(\cdot, \cdot)_a$ to \mathfrak{a} , has the same matrix B_a in the basis Π_a . In particular, we have that :

$$(\rho_1(\alpha'), \rho_1(\alpha'))_a = 2[(n_1 + 2n_3)^2 + (n_2 + 2n_3)^2 - 3(n_1 + 2n_3)(n_2 + 2n_3)],$$

since $\rho_1(\alpha') = (n_1 + 2n_3)\gamma_1 + (n_2 + 2n_3)\gamma_2$.

Using (5.1), it is not difficult to check that

$$(\rho_1(\alpha'), \rho_1(\alpha'))_a = 2[(\alpha', \alpha')_1 - (n_3 - n_4)^2 - 5n_3^2 - n_4^2] \tag{5.2}$$

Suppose $n_3 = 0$, then, since $\text{supp}(\alpha')$ is connected, we have that $\alpha' = n_1\alpha'_1 + n_2\alpha'_2$ or $\alpha' = \alpha'_4$. Hence $\rho_1(\alpha') = n_1\gamma_1 + n_2\gamma_2 \in \Sigma$ or $\rho_1(\alpha') = 0$.

Suppose $n_3 \neq 0$, then, since $(\alpha', \alpha')_1 \leq 2$, one can see, using (5.2), that

$$(\rho_1(\alpha'), \rho_1(\alpha'))_a < 0.$$

As Σ is hyperbolic and $\rho_1(\alpha') \in \mathbb{N}\gamma_1 + \mathbb{N}\gamma_2$, we deduce that $\rho_1(\alpha')$ is a positive imaginary root of Σ (see [12, 5.10]). It follows that $\rho_1(\Delta'^+) = \Sigma^+ \cup \{0\}$.

To see that \mathfrak{m}^1 is finitely Σ -graded, it suffices to prove that, for $\gamma = m_1\gamma_1 + m_2\gamma_2 \in \Sigma^+ \cup \{0\}$, the set $\{\alpha' \in \Delta'^+, \rho_1(\alpha') = \gamma\}$ is finite. Note that if $\alpha' = \sum_{i=1}^4 n_i\alpha'_i \in \Delta'^+$ satisfying $\rho_1(\alpha') = \gamma$, then $n_i + 2n_3 = m_i$, $i = 1, 2$. In particular, there are only finitely many possibilities for n_i , $i = 1, 2, 3$. The same argument as the one used in the proof of Proposition 2.13 shows also that there are only finitely many possibilities for n_4 .

3) Recall that $\mathfrak{m} \subset \mathfrak{m}^1 \subset \mathfrak{g}$. The fact that \mathfrak{g} is finitely Δ' -graded with grading subalgebra \mathfrak{m}^1 and \mathfrak{m}^1 is finitely Σ -graded implies that \mathfrak{g} is finitely Σ -graded (cf. Lemma 1.5). Let $I = \{1, 2, 3, 4, 5, 6\}$, then the root basis Π_a of Σ is adapted to the root basis Π of Δ and we have $I_{re} = \{1, 2, 5, 6\}$ (not connected), $\Gamma_1 = \{1, 5\}$, $\Gamma_2 = \{2, 6\}$, $J = \{4\}$, $J_{re} = \emptyset$, $I'_{im} = \{3\}$ and $J^\circ = J = \{4\}$. Note that, for this example, $\mathfrak{g}(I_{re})$, which is Σ -graded, is isomorphic to $\mathfrak{m} \times \mathfrak{m}$. This gradation corresponds to that of the pseudo-complex real form of $\mathfrak{m} \times \mathfrak{m}$ (i.e. the complex Kac-Moody algebra \mathfrak{m} viewed as real Lie algebra) by its restricted reduced root system. Since the pair $(I_3, J_3) = (\{3, 4\}, \{4\})$ is not admissible, it is not possible to build a Kac-Moody algebra \mathfrak{g}^J grading finitely \mathfrak{g} and maximally finitely Σ -graded.

Acknowledgments. We thank the anonymous referee for his/her valuable comments and suggestions.

References

- [1] Allison, B., G. Benkart and Y. Gao, *Central extensions of Lie algebras graded by finite root systems*, Math. Ann. **316** (2000), 499–527.
- [2] Back-Valente, V., N. Bardy-Panse, H. Ben Messaoud and G. Rousseau, *Formes presque déployées d’algèbres de Kac-Moody, Classification et racines relatives*, J. of Algebra **171** (1995), 43–96.
- [3] Bardy-Panse, N., «Systèmes de racine infinis», Mémoire de la S.M.F **65**, 1996.
- [4] Bardy, N., *Sous-algèbres birégulières d’une algèbre de Kac-Moody-Borcherds*, Nagoya Math. J. **156** (1999), 1–83.
- [5] Benkart, G., and E. Zelmanov, *Lie algebras graded by finite root systems and intersection matrix algebras*, Invent. Math. **126** (1996), 1–45.
- [6] Ben Messaoud, H., *Almost split real forms for hyperbolic Kac-Moody Lie algebras*, J. Phys. A. Math. Gen **39** (2006), 13659–13690.
- [7] Ben Messaoud, H., and N. Fradi, *Imaginary gradations of Kac-Moody Lie algebras*, In preparation.
- [8] Berman, S. and R. Moody, *Lie algebras graded by finite root systems*, Invent. Math. **108** (1992), 323–347.
- [9] Borcherds, R. E., *Generalized Kac-Moody algebras*, J. of Algebra **115** (1988), 501–512.
- [10] Bourbaki, N., «Groupes et algèbres de Lie», Chap 4, 5 et 6», Masson, Paris, 1981.
- [11] Carter, R., “Lie algebras of finite and affine type,” Cambridge University Press, 2005.
- [12] Kac, V. G., “Infinite dimensional Lie algebras,” Third Edition, Cambridge University Press, 1990.
- [13] Kac, V. G., and S. P. Wang, *On automorphisms of Kac-Moody algebras and groups*, Advances in Math. **92** (1992), 129–195.
- [14] Moody, R. V., *Root systems of hyperbolic type*, Advances in Mathematics **33** (1979), 144–160.
- [15] Neher, E., *Lie algebras graded by J -graded root systems and Jordan pairs covered by grids*, Amer. J. Math. **118** (1996), 439–491.
- [16] Nervi, J., *Algèbres de Lie simples graduées par un système de racines et sous-algèbres C -admissibles*, J. of Algebra **223** (2000), 307–343.
- [17] —, *Affine Kac-Moody algebras graded by affine root systems*, J. of Algebra **253** (2002), 50–99.

- [18] Peterson, D. H., and V. G. Kac, *Infinite flag varieties and conjugacy theorems*, Proc. Natl. Acad. Sc. USA **80** (1983), 1778–1782.
- [19] Rémy, B., «Groupes de Kac-Moody déployés et presque déployés», *Astérisque* **277**, 2002.
- [20] Rousseau, G., *Groupes de Kac-Moody déployés sur un corps local, II Mesures ordonnées*, ArXiv 1009.0135v2.
- [21] Rubenthaler, H., *Construction de certaines sous-algèbres remarquables dans les algèbres de Lie semi-simples*, J. of Algebra **81** (1983), 268–278.
- [22] Tits, J., *Uniqueness and presentation of Kac-Moody groups over fields*, J. of Algebra **105** (1987), 542–573.

Hechmi Ben Messaoud
Université de Monastir
Faculté des Sciences
Département de Mathématiques
5019 Monastir. Tunisie
hechmi.benmessaoud@fsm.rnu.tn

Guy Rousseau
Université de Lorraine
CNRS, UMR 7502
Institut Élie Cartan de Lorraine
Vandœuvre-lès-Nancy, 54506. France
guy.rousseau@univ-lorraine.fr

Received July 22, 2013
and in final form September 4, 2013